

MIRROR SYMMETRY: LECTURE 17

DENIS AUROUX
NOTES BY KARTIK VENKATRAM

1. COHERENT SHEAVES ON A COMPLEX MANIFOLD (CONTD.)

We now recall the following definitions from category theory.

Definition 1. An additive category is one in which $\text{Hom}(A, B)$ are abelian groups, composition is distributive, and there is a direct sum \oplus and a zero object 0. An abelian category is an additive category s.t. every morphism has a kernel and cokernel, e.g. a kernel of $f : A \rightarrow B$ is a morphism $K \rightarrow A$ s.t. $g : C \rightarrow A$ factors through K uniquely iff $f \circ g = 0$.

One can define complexes in an additive category, but one needs to be in an abelian category to have notions of exact sequences and cohomology. Recall that, given chain complexes C_*, D_* , a chain map $f : C_* \rightarrow D_*$ is a collection of maps $f_i : C_i \rightarrow D_i$ commuting with δ . Given two such maps $f = \{f_i\}, g = \{g_i\}$, we call them *homotopic* if there is a map $h : A \rightarrow B[-1]$ (B shifted down by 1) s.t. $f - g = d_B h + h d_A$, i.e.

$$(1) \quad \begin{array}{ccccccc} \cdots & \longrightarrow & A_{i-1} & \xrightarrow{d_{i-1}} & A_i & \xrightarrow{d_i} & A_{i+1} \longrightarrow \cdots \\ & & \downarrow f_{i-1} & \swarrow h_{i-1} & \downarrow f_i & \swarrow h_i & \downarrow f_{i+1} \\ & & B_{i-1} & \xrightarrow{d_{i-1}} & B_i & \xrightarrow{d_i} & B_{i+1} \longrightarrow \cdots \end{array}$$

A chain map is a *quasi-isomorphism* if the induced maps on cohomology are isomorphisms. This is stronger than $H^*(C_*) \cong H^*(D_*)$. For \mathcal{A} an abelian category, the category of bounded chain complexes is the differential graded category whose objects are bounded chain complexes in \mathcal{A} and whose morphisms are “pre-homomorphisms” of complexes $\text{Hom}^k(A_*, B_*) = \bigoplus_i \text{Hom}_{\mathcal{A}}(A_i, B_{i+k})$: it is equipped with a differential δ where

$$(2) \quad f \in \text{Hom}^k(A_*, B_*) \implies \delta(f) = d_B f + (-1)^{k+1} f d_A \in \text{Hom}^{k+1}(A_*, B_*)$$

Chain maps are precisely the elements of $\text{Ker}(\delta : \text{Hom}^0 \rightarrow \text{Hom}^1)$, and the nullhomotopic maps are elements of $\text{im}(\delta : \text{Hom}^{-1} \rightarrow \text{Hom}^0)$, so $H^0 \text{Hom}(A, B)$ gives the space of chain maps up to homotopy.

Definition 2. For \mathcal{A} an abelian category, the bounded derived category $D^b(\mathcal{A})$ is the triangulated category whose objects are bounded chain complexes in \mathcal{A} and

whose morphisms are given by chain maps up to homotopy localizing w.r.t. quasi-isomorphisms. That is, quasi-isomorphisms are formally inverted; for any quasi-isomorphism s , we add a morphism s^{-1} . More precisely, $\text{Hom}_{D^b(\mathcal{A})}(A_*, B_*) = \{A \xleftarrow{s} A' \xrightarrow{f} B\} / \sim$ where s is a quasi-isomorphism, f is a chain map, and \sim is homotopy equivalence. We similarly define the categories $D^+(\mathcal{A}), D^-(\mathcal{A})$ of chain complexes bounded above/below.

To explain the notion of triangulated category, recall the following:

- In the category of topological spaces (or simplicial complexes), there are no kernels and cokernels. Given a map f , however, the mapping cone $C_f = (X \times [0, 1]) \sqcup Y / (x, 0) \sim (x', 0), (x, 1) \sim f(x)$ acts as both simultaneously. There are natural maps $i : Y \rightarrow C_f$ (inclusion) and $q : C_f \rightarrow \Sigma X$ (collapsing Y), and we obtain a sequence of topological spaces

$$(3) \quad X \xrightarrow{f} Y \xrightarrow{i} C_f \xrightarrow{q} \Sigma X \rightarrow \dots$$

with compositions null-homotopic. This gives a long exact sequence of

$$(4) \quad H_i(X) \rightarrow H_i(Y) \rightarrow H_i(C_f) \rightarrow H_i(\Sigma X) = H_{i-1}(X) \rightarrow H_i(\Sigma Y) = H_{i-1}(Y)$$

- If X, Y are simplicial complexes, f a simplicial map, C_f defined analogously is a simplicial complex, with i -cells given by cones on $(i-1)$ -cells of X and i -cells of Y . The boundary map is given by the matrix $\begin{pmatrix} \partial_X & 0 \\ f & \partial_Y \end{pmatrix}$.
- If A^* and B^* are complexes, f a chain map, we define $C_f = A[1] \oplus B$, i.e. $C_f^i = A^{i+1} \oplus B^i$. The boundary map is $\delta = \begin{pmatrix} \delta_A[1] & 0 \\ f & \delta_B \end{pmatrix}$. Note that, if A, B are single objects, $\text{Cone}(f : A \rightarrow B)$ is just $\{0 \rightarrow A \xrightarrow{f} B \rightarrow 0\}$. We have natural chain maps $B^* \xrightarrow{i} C_f^*$ (subcomplex) and $C_f^* \xrightarrow{q} A^*[1]$ (quotient complex). As before, $A^*[1]$ is quasi-isomorphic to $\text{Cone}(i : B^* \rightarrow C_f^*)$.
- Finally, in the derived category, the inversion of quasi-isomorphisms gives us *exact triangles*

$$(5) \quad \begin{array}{ccc} A^* & \xrightarrow{\quad} & B^* \\ & \swarrow \quad \searrow & \\ & [1] \quad C^* & \end{array}$$

with

$$(6) \quad H^i(A) \rightarrow H^i(B) \rightarrow H^i(C) \rightarrow H^{i+1}(A) \rightarrow \dots$$

Definition 3. A triangulated category is an additive category with a shift functor $[1]$ and a set of distinguished triangles satisfying various axioms:

- $\forall X, X \xrightarrow{\text{id}} X \rightarrow 0 \rightarrow X[1]$ is distinguished,
- $\forall X \rightarrow Y$, there is a distinguished triangle $X \xrightarrow{u} Y \rightarrow Z \rightarrow X[1]$ (Z is called the mapping cone of f).
- The rotation of any distinguished triangle is distinguished, i.e. for $X \rightarrow Y \rightarrow Z \rightarrow X[1]$ distinguished, $Y \rightarrow Z \rightarrow X[1] \rightarrow Y[1]$ and $Z \rightarrow X[1] \rightarrow Y[1] \rightarrow Z[1]$ are distinguished.
- Given a square

$$(7) \quad \begin{array}{ccc} X & \xrightarrow{f} & Y \\ \downarrow & & \downarrow \\ X' & \xrightarrow{f'} & Y' \end{array}$$

there is a map between the mapping cones of f, f' that makes everything commute in the induced map of distinguished triangles

$$(8) \quad \begin{array}{ccccccc} X & \longrightarrow & Y & \longrightarrow & Z & \longrightarrow & X[1] \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ X' & \longrightarrow & Y' & \longrightarrow & Z' & \longrightarrow & X'[1] \end{array}$$

- Given a pair of maps $X \xrightarrow{u} Y \xrightarrow{v} Z$, there are maps between the mapping cones $C_u, C_v, C_{v \circ u}$ of u, v , and $v \circ u$ that make every commute in the induced maps of distinguished triangles.

$$(9) \quad \begin{array}{ccccc} & & C_{u \circ v} & & \\ & \swarrow \text{dashed} & \uparrow \text{dashed} & \searrow \text{dashed} & \\ C_u & \xleftarrow{[1]} & & \xrightarrow{[1]} & C_v \\ \downarrow [1] & & \downarrow [1] & & \downarrow [1] \\ X & \xrightarrow{v \circ u} & [1] & \longrightarrow & Z \\ \downarrow u & & & & \downarrow v \\ & & Y & & \end{array}$$

1.1. Derived functors. Let $F : \mathcal{A} \rightarrow \mathcal{B}$ be a left exact functor between abelian categories. $\mathcal{R} \subset \mathcal{A}$ is called an *adapted class of objects* for F if

- \mathcal{R} is stable under direct sums,
- for C^* an acyclic complex of objects in \mathcal{R} , $F(C^*)$ is acyclic, and
- $\forall A \in \mathcal{A}, \exists R \in \mathcal{R}$ s.t. $0 \rightarrow A \xrightarrow{i} R$.

For instance, the set of injective objects is such an adapted class. Let $K^+(\mathcal{R})$ be the homotopy category of complexes bounded below of objects in \mathcal{R} . RF gives a composition $D^+(\mathcal{A}) \rightarrow K^+(\mathcal{R}) \xrightarrow{F} D^+(\mathcal{B})$, where the first map is induced by resolution by objects of R . The map $D^+(\mathcal{A}) \rightarrow D^+(\mathcal{B})$ is exact, i.e. it maps exact triangles to exact triangles, and $R^i F = H^i(RF)$.

1.2. Extensions. Let $A, B \in \mathcal{A} \hookrightarrow D^b(\mathcal{A})$ be single object complexes concentrated in degree 0, so $B[k]$ is concentrated in degree $-k$.

Proposition 1. $\mathrm{Hom}_{D^b(\mathcal{A})}(A, B[k]) = \mathrm{Ext}_{\mathcal{A}}^k(A, B)$.

We can use this to define a product $\mathrm{Ext}_{\mathcal{A}}^k(A, B) \otimes \mathrm{Ext}_{\mathcal{A}}^\ell(B, C) \rightarrow \mathrm{Ext}_{\mathcal{A}}^{k+\ell}(A, C)$ as a composition $A \rightarrow B[k] \rightarrow C[k+\ell]$ in $D^b(\mathcal{A})$.

Example. For $k = 1$, we have

$$(10) \quad \begin{array}{ccccccc} 0 & \longrightarrow & 0 & \longrightarrow & A & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & B & \longrightarrow & 0 & \longrightarrow & 0 \end{array}$$

There are no chain maps, but we can invert quasi-isomorphisms. If we have an extension $0 \rightarrow A \xrightarrow{f} B \xrightarrow{g} C \rightarrow 0$ in \mathcal{A} , we have chain maps

$$(11) \quad \begin{array}{ccccccc} 0 & \longrightarrow & 0 & \longrightarrow & C & \longrightarrow & 0 \\ & & & & \uparrow g & & \\ 0 & \longrightarrow & A & \xrightarrow{f} & B & \longrightarrow & 0 \\ & & \downarrow \mathrm{id} & & & & \\ 0 & \longrightarrow & A & \longrightarrow & 0 & \longrightarrow & 0 \end{array}$$

giving an element in $\mathrm{Hom}_{D^b(\mathcal{A})}(C, A[1]) = \mathrm{Ext}^1(C, A)$.

There are two ways to understand the above proposition. First, if \mathcal{A} has enough injectives, take a resolution of B by a complex $I^0 \rightarrow I^1 \rightarrow \dots$ quasi-isomorphic to B : the chain maps from A to I^* are, up to homotopy, isomorphic to $H^k(\mathrm{Hom}(A, I^*)) \cong \mathrm{Ext}^k(A, B)$. Second, we can check the definition of a derived functor. Given a short exact sequence $0 \rightarrow A \xrightarrow{f} B \xrightarrow{g} C \rightarrow 0$ in \mathcal{A} , we get an exact triangle $A \xrightarrow{f} B \xrightarrow{g} C \xrightarrow{w} A[1]$ quasi-isomorphic to a distinguished triangle with $\mathrm{Cone}(f)$.

Proposition 2. *For an exact triangle $A \xrightarrow{f} B \xrightarrow{g} C \xrightarrow{h} A[1]$ and an object E , we have long exact sequences*

(12)

$$\begin{aligned} \cdots \rightarrow \mathrm{Hom}(E, A[i]) \xrightarrow{f_*} \mathrm{Hom}(E, B[i]) \xrightarrow{g_*} \mathrm{Hom}(E, C[i]) \xrightarrow{h_*} \mathrm{Hom}(E, A[i+1]) \rightarrow \cdots \\ \cdots \rightarrow \mathrm{Hom}(A[i+1], E) \xrightarrow{h^*} \mathrm{Hom}(C[i], E) \xrightarrow{g^*} \mathrm{Hom}(B[i], E) \xrightarrow{f^*} \mathrm{Hom}(A[i], E) \rightarrow \cdots \end{aligned}$$