

# MIRROR SYMMETRY: LECTURE 18

DENIS AUROUX  
NOTES BY KARTIK VENKATRAM

## 1. DERIVED FUKAYA CATEGORY

Last time: derived categories for abelian categories (e.g.  $D^b\text{Coh}(X)$ ). This time: the derived Fukaya category. We start with an  $A_\infty$ -category  $\mathcal{A}$  and obtain a triangulated category via “twisted complexes”. Recall that in an  $A_\infty$ -category,  $\text{hom}_{\mathcal{A}}(X, Y)$  is a graded vector space equipped with maps

$$(1) \quad m_k : \text{hom}_{\mathcal{A}}(X_0, X_1) \otimes \cdots \otimes \text{hom}_{\mathcal{A}}(X_{k-1}, X_k) \rightarrow \text{hom}_{\mathcal{A}}(X_0, X_k)[2 - k]$$

1) Additive enlargement: we define the category  $\Sigma\mathcal{A}$  to be the category whose objects are finite sums  $\bigoplus X_i[k_i]$ ,  $X_i \in \mathcal{A}$ ,  $k_i \in \mathbb{Z}$  and whose maps are

$$(2) \quad \text{hom}_{\Sigma\mathcal{A}}(\bigoplus_i X_i[k_i], \bigoplus_j Y_j[\ell_j]) = \bigoplus_{i,j} \text{hom}_{\mathcal{A}}(X_i, Y_j)[\ell_j - k_i]$$

Note that we have induced multiplication maps

$$(3) \quad m_k(a_k, \dots, a_1)^{ij} = \sum_{i_1, \dots, i_{k-1}} m_k(a_k^{i_{k-1}, j}, \dots, a_1^{i_1, j})$$

2) Twisted complexes: we define the category  $\text{Tw}\mathcal{A}$  to be the category whose objects are twisted complexes  $(X, \delta_X)$ ,

$$(4) \quad X = \bigoplus_i X_i[k_i] \in \Sigma\mathcal{A}, \delta_X = (\delta_X^{ij}) \in \text{hom}_{\Sigma\mathcal{A}}^1(X, X)$$

(i.e.  $\delta_X$  a degree 1 endomorphism) s.t.

- $\delta_X$  is strictly lower-triangular, and
- $\sum_{k=1}^{\infty} m_k(\delta_x, \dots, \delta_x) = 0$ . It is a finite sum because  $\delta_X$  is lower triangular, and generalizes  $\delta_X \circ \delta_X = 0$ .

*Example.* For a simple map  $f : X_1 \rightarrow X_2$ ,  $f \in \text{hom}_{\mathcal{A}}^1(X_1, X_2)$ , the condition is  $m_1(f) = 0$ . Now, for maps  $X_1[2] \xrightarrow{f} X_2[1] \xrightarrow{g} X_3$  and  $X_1[2] \xrightarrow{h} X_3$ ,

$$(5) \quad \begin{aligned} g &\in \text{hom}^0(X_2, X_3) = \text{hom}^1(X_2[1], X_3) \\ f &\in \text{hom}^0(X_1[1], X_2[1]) = \text{hom}^1(X_1[2], X_2[1]) \\ h &\in \text{hom}^{-1}(X_1, X_3) = \text{hom}^1(X_1[2], X_3) \end{aligned}$$

the condition is  $m_1(f) = m_1(g) = 0$  and  $m_2(g, f) + m_1(h) = 0$ .

The morphisms in the category of twisted complexes are

$$(6) \quad \text{hom}_{\text{Tw}\mathcal{A}}((X, \delta_X), (Y, \delta_Y)) = \text{hom}_{\Sigma\mathcal{A}}(X, Y)$$

and

$$(7) \quad m_k^{\text{Tw}\mathcal{A}}(a_k, \dots, a_1) = \sum_{i_0, \dots, i_k} \pm m_{k+i_0+\dots+i_k}^{\Sigma\mathcal{A}} \left( \underbrace{\delta_{X_k}, \dots, \delta_{X_k}}_{i_k}, a_k, \underbrace{\delta_{X_{k-1}}, \dots, \delta_{X_{k-1}}}_{i_{k-1}}, \dots, \underbrace{\delta_{X_1}, \dots, \delta_{X_1}}_{i_1}, a_1, \underbrace{\delta_{X_0}, \dots, \delta_{X_0}}_{i_0} \right)$$

$\text{Tw}\mathcal{A}$  is a *triangulated*  $A_\infty$ -category, i.e. there are mapping cones satisfying the usual axioms.

*Example.* For  $a \in \text{hom}(X, Y)$ ,

$$(8) \quad m_1^{\text{Tw}\mathcal{A}}(a) = m_1(a) \pm m_2(\delta_Y, a) \pm m_2(a, \delta_X) + \text{higher terms}$$

This is a generalization of being a chain map up to homotopy.

3) We now take the cohomology category  $D(\mathcal{A}) := H^0(\text{Tw}\mathcal{A})$ , which is an honest triangulated category. The objects of the two categories are the same, but now our morphisms are  $\text{hom}^{D(\mathcal{A})}(X, Y) := H^0(\text{hom}^{\text{Tw}\mathcal{A}}(X, Y), m_1^{\text{Tw}(\mathcal{A})})$ . Note that  $\text{hom}^{D(\mathcal{A})}(X, Y[k]) = H^k(\text{hom}^{\text{Tw}\mathcal{A}}(X, Y), m_1^{\text{Tw}\mathcal{A}})$ . The composition is induced by  $m_2^{\text{Tw}\mathcal{A}}$  on cohomology.

*Remark.* There is a variant of this called a *split-closed derived category*. Let  $\mathcal{A}$  be a linear category,  $X \in \mathcal{A}, p \in \text{hom}_{\mathcal{A}}(X, X)$  s.t.  $p^2 = p$  (idempotent). Define the image of  $p$  to be an object  $Y$ , and add maps  $u : X \rightarrow Y, v : Y \rightarrow X$  s.t.  $u \circ v = \text{id}_Y, v \circ u = p$ . That is, we enlarge  $\mathcal{A}$  to add these objects and maps, and define the split closure to be the category whose objects are  $(X, p)$  with  $p$  idempotent, and morphisms  $\text{hom}((X, p), (Y, p')) = p' \text{hom}(X, Y)p$ . This is more complicated in the  $A_\infty$  setting.

Geometrically, some exact triangles in  $DFuk(M)$  are given by Lagrangian connected sums (FOOO) and Dehn twists (Seidel).

- For an example of the latter, given a cylinder with a Lagrangian circle  $S$ , we can obtain a symplectomorphism  $\tau_S \in \text{Symp}(M, \omega)$  which is the identity outside a neighborhood of  $S$  and, within that neighborhood, twists the cylinder around (in higher dimensions, define this using the geodesic flow in a neighborhood of  $S \cong T^*S$ ). If  $L$  is Lagrangian, then  $\tau_S(L)$  is Lagrangian as well. By Seidel, there exists an exact triangle in

$DFuk(M)$ :

$$(9) \quad \begin{array}{ccc} HF^*(S, L) \otimes S & \xrightarrow{t} & L \\ & \nwarrow \quad \nearrow & \\ & [1] \quad \tau_S(L) & \end{array}$$

These correspond to long exact sequences for  $HF(L', -)$ .

- In the former situation, for  $L_1, L_2$  (graded) Lagrangians,  $L_1 \cap L_2 = \{p\}$  of index 0, we can construct the connected sum  $L_1 \#_p L_2$ , which looks locally like  $\tau_{L_1}(L_2)$  if  $L_1$  is a sphere and is given by  $\text{Cone}(L_1 \xrightarrow{p} L_2)$  in general (consider this vs. “ $L_1[1] \cup_p L_2 \simeq \text{Cone}(L_1 \xrightarrow{0} L_2)$ ”). For instance, in the torus  $T^2$ , consider two independent loops  $\alpha$  of degree 2 and  $\beta$  of degree 1, with two points of intersection  $p, q$ . Then  $\text{Cone}(\alpha \xrightarrow{p+q} \beta) \simeq \gamma_1 \oplus \gamma_2$  is disconnected, where  $\gamma_1, \gamma_2$  are degree 1 loops. If we only started with  $\alpha, \beta$ , the triangulated envelope contains  $\gamma_1 \oplus \gamma_2$ , but not  $\gamma_1, \gamma_2$  separately. The split-closure does contain them.
- Now, if we start with two independent generators of the torus, successive Dehn twists give all the homotopy classes of loops in  $T^2$ , but each homotopy class contains infinitely many non-Hamiltonian isotopic Lagrangians. To generate  $DFuk(T^2)$  as a triangulated envelope, we need (for instance) one horizontal loop and infinitely many vertical loops. On the other hand,  $\alpha, \beta$  above are split generators. The key point is that  $\text{Cone}(\alpha \xrightarrow{p+T^q q} \beta)$  gives deformed loops, direct sums of which vary continuously within a homotopy class. But many cones and idempotents have no obvious geometric interpretation. For instance, the Clifford torus  $T = \{|x| = |y| = |z|\} \subset \mathbb{CP}^2$  has idempotents in  $HF(T, T)$  without any obvious geometric interpretation.