


partition λ of $n \geq 0$:

$$\lambda = (\lambda_1, \lambda_2, \dots)$$


$$\lambda_1 \geq \lambda_2 \geq \dots \geq 0$$

$$\sum \lambda_i = n$$

Young diagram of $\lambda = (4, 4, 1)$:



standard Young tableau (SYT) of shape $(4, 4, 1)$:




$f^\lambda = \#$ of SYT of shape λ

1 2 3 4	1 2 3 5	1 2 3 6
5 6	4 6	4 5

1 2 4 5	1 2 4 6	1 2 5 6
3 6	3 5	3 4

1 3 4 5	1 3 4 6	1 3 5 6
2 6	2 5	2 4

$$f^{4,2} = 9$$



$H(u)$: **hook** at (or of) u

$h(u) = \#H(u)$: **hook length** at u

8	7	5	4	1
6	5	3	2	
5	4	2	1	
2	1			

Frame-Robinson-Thrall hook length formula (1954):

$$f^\lambda = \frac{n!}{\prod_{u \in \lambda} h(u)}$$

hook lengths : 5 4 2 1
 2 1

$$f^{4,2} = \frac{6!}{5 \cdot 4 \cdot 2 \cdot 2 \cdot 1 \cdot 1} = 9$$

“nice” bijective proof by Novelli-Pak-Stoyanovskii (1997)

Robinson-Schensted-Knuth (RSK)

algorithm: $w \xrightarrow{\text{rsk}} (P, Q)$, where $w \in \mathfrak{S}_n$ and P, Q are SYT of same shape $\lambda \vdash n$

Note. Schensted = Ea Ea

(ea.ea.home.mindspring.com)

$$w = 4273615$$

4 1

2 1

4 2

2**7** 13

4 2

2**3** 13

4**7** 24

23**6** 135

47 24

136 135

27 24

4 6

13**5** 135

2**6** 24

4**7** 67

An element j bumps the smallest $i > j$.

First symmetry property.

$$w = 4273615 \xrightarrow{\text{rsk}} \begin{array}{cc} 135 & 135 \\ 26 & 24 \\ 47 & 67 \end{array}$$

$$w^{-1} = 6241753 \xrightarrow{\text{rsk}} \begin{array}{cc} 135 & 135 \\ 24 & 26 \\ 67 & 47 \end{array}$$

Theorem (Schützenberger) *If $w \xrightarrow{\text{rsk}} (P, Q)$, then*

$$w^{-1} \xrightarrow{\text{rsk}} (Q, P).$$

Corollary. *Let $t(n)$ denote the number of SYT with n squares. Then*

$$t(n) = \#\{w \in \mathfrak{S}_n : w^2 = 1\}.$$

$$w = 318496725$$

is(w) := length of longest increasing
subsequence of $w \in \mathfrak{S}_n$

$$\text{is}(318496725) = 4$$

ds(w) := length of longest decreasing
subsequence of $w \in \mathfrak{S}_n$

$$\text{ds}(318496725) = 3$$

$$\begin{array}{ccc}
& & 1\ 2\ 5\ 7 & 1\ 3\ 5\ 7 \\
318496725 & \xrightarrow{\text{rsk}} & 3\ 4\ 6 & 2\ 4\ 6 \\
& & 8\ 9 & 8\ 9
\end{array}$$

Theorem. Let $w \xrightarrow{\text{rsk}} (P, Q)$,
 $\text{shape}(P) = (\lambda_1, \lambda_2, \dots)$.

Then

$$\text{is}(w) = \lambda_1.$$

Proof (sketch). Let the first row be

$$b_1, b_2, \dots, b_k.$$

Straightforward to show by induction on n that b_i is the rightmost element j of w for which the longest increasing subsequence of w ending at j has length i . \square

Second symmetry property.

$$w = a_1 a_2 \cdots a_n, \quad \bar{w} := a_n \cdots a_2 a_1$$

Theorem (Schensted). *If $w \xrightarrow{\text{rsk}} (P, Q)$, then*

$$\bar{w} \xrightarrow{\text{rsk}} (P^t, \text{evac}(Q)^t).$$

Corollary. *Let $w \xrightarrow{\text{rsk}} (P, Q)$, $\text{shape}(P) = (\lambda_1, \lambda_2, \dots)$. Then*

$$\text{ds}(w) = \lambda'_1 = \ell(\lambda).$$

Corollary (Erdős-Szekeres). *Let $w \in \mathfrak{S}_{pq+1}$. Then either*

$$\text{is}(w) > p \text{ or } \text{ds}(w) > q.$$


Proof. Let

$$w \xrightarrow{\text{rsk}} (P, Q), \quad \text{shape}(P) = \lambda.$$

$$\text{is}(w) \leq p \Rightarrow \lambda_1 \leq p$$

$$\text{ds}(w) \leq q \Rightarrow \lambda'_1 \leq q$$

$$\Rightarrow |\lambda| \leq pq.$$



Corollary. *Let $p \leq q$ (say). Then*

$$\begin{aligned} \#\{w \in \mathfrak{S}_{pq} : \text{is}(w) = p, \text{ds}(w) = q\} \\ = (f^{q \times p})^2, \end{aligned}$$

where $f^{q \times p} =$

$$(pq)!$$

$$1^1 2^2 \cdots p^p (p+1)^p \cdots q^p (q+1)^{p-1} \cdots (p+q-1)^1.$$

7	6	5	4	3
6	5	4	3	2
5	4	3	2	1

A **reverse semistandard tableau** T
of shape $(5, 4, 3)$:

$$\begin{array}{cccccc} 6 & 6 & 4 & 2 & 2 & \\ & 4 & 4 & 3 & 1 & \\ & & 2 & 1 & 1 & \end{array}$$

(weakly decreasing in rows, strictly decreasing in columns)

$$A = (a_{ij}) = \begin{bmatrix} 0 & 0 & 0 & 2 \\ 0 & 2 & 1 & 0 \\ 1 & 0 & 1 & 0 \end{bmatrix}$$

$$w_A = \begin{pmatrix} 3 & 3 & 2 & 2 & 2 & 1 & 1 \\ 3 & 1 & 3 & 2 & 2 & 4 & 4 \end{pmatrix}$$

3 3

3**1** 33

3**3** 33
1 2

33**2** 332
 1 2

332**2** 3322
 1 2

4322 3322
3 2
1 1

4**4**22 3322
 3**3** 21
 1 1

- a_{ij} columns of w_A are equal to $\begin{smallmatrix} i \\ j \end{smallmatrix}$.
- Top row of w_A is weakly decreasing.
- Bottom row is weakly decreasing under equal elements of the top row.
- An element j bumps the largest element $i < j$.

Lemma (simple). Let $A \xrightarrow{\text{rsk}} (P, Q)$. Equal elements of Q are inserted left-to-right, allowing the construction of the inverse map $(P, Q) \rightarrow A$.

Note.

j appears in P : $\sum_i a_{ij}$ times (j th column sum)

i appears in Q : $\sum_j a_{ij}$ times (i th row sum)

$$\sum (\text{entries of } P) = \sum_{i,j} j a_{ij}$$

$$\sum (\text{entries of } Q) = \sum_{i,j} i a_{ij}$$

$$\#(\text{entries of } P \text{ or } Q) = \sum_{i,j} a_{ij}$$

Plane partition of $n \geq 0$:

$$\pi = (\pi_{ij})_{i,j \geq 1}$$

$$\pi_{ij} \geq \pi_{i+1,j}, \quad \pi_{ij} \geq \pi_{i,j+1}, \quad \sum \pi_{ij} = n$$

$a(n)$ = # plane partitions of n

553221

53311

52211

221

3 21 111 2 1 11

 1 1 1

 1

$$a(3) = 6$$

MacMahon:

$$\sum_{n \geq 0} a(n)x^n = \prod_{i \geq 1} (1 - x^i)^{-i}.$$

Compare Euler's theorem for $p(n)$, the number of partitions (or one-row plane partitions) of n :

$$\sum_{n \geq 0} p(n)x^n = \prod_{i \geq 1} (1 - x^i)^{-1}$$

(much easier).


$$A = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 2 \\ 1 & 0 & 1 \\ 0 & 2 & 0 \end{bmatrix}$$

$$\xrightarrow{\text{rsk}} \begin{array}{cc} 33332 & 44321 \\ 222 & 322 \\ 1 & 1 \end{array}$$

$$\xrightarrow{\text{merge}} \begin{array}{c} 33332 \\ 3331 \\ 332 \\ 21 \end{array} = \pi_A$$

Merge column-by-column.

merge of $(5, 4, 2)$ and $(6, 3, 1)$ is $(5, 5, 4, 2, 1, 1)$:



$$\begin{aligned}
|\pi_A| &= \sum (\text{entries of } P) \\
&\quad + \sum (\text{entries of } Q) \\
&\quad - \#(\text{entries of } P \text{ or } Q) \\
&= \sum_{i,j} (i + j - 1) a_{ij}
\end{aligned}$$

$$\begin{aligned}
\Rightarrow \sum_{\pi} x^{|\pi|} &= \sum_A x^{\sum (i+j-1) a_{ij}} \\
&= \prod_{i,j \geq 1} \left(\sum_{a_{ij} \geq 0} x^{(i+j-1) a_{ij}} \right) \\
&= \prod_{i,j \geq 1} (1 - x^{i+j-1})^{-1} \\
&= \prod_{i \geq 1} (1 - x^i)^{-i}
\end{aligned}$$


slight modification \Rightarrow

$$\sum_{\substack{\pi \\ \leq r \text{ rows} \\ \leq s \text{ cols}}} x^{|\pi|} = \prod_{i=1}^r \prod_{j=1}^s (1 - x^{i+j-1})^{-1}$$

More difficult (MacMahon):

$$\sum_{\substack{\pi \\ \leq r \text{ rows} \\ \leq s \text{ cols} \\ \max \leq t}} x^{|\pi|} = \prod_{i=1}^r \prod_{j=1}^s \prod_{k=1}^t \frac{1 - x^{i+j+k-1}}{1 - x^{i+j+k-2}}$$


Oscillating tableaux: start with \emptyset , add or remove a square at each step.



shape $(3, 1)$, length 8

$$\tilde{f}_n^\lambda = \#\{\text{osc. tab. of shape } \lambda, \text{ length } n\}$$

M : (complete) **matching** on $1, 2, \dots, 2n$



$$\Phi(M) = (\emptyset \square \begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array} \square \square \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array} \begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array} \square \emptyset)$$

Φ is a bijection between matchings on $1, 2, \dots, 2n$ and oscillating tableaux of length $2n$, shape \emptyset .

Hence

$$\begin{aligned} \tilde{f}_{2n}^{\emptyset} &= \sum_{\lambda} \left(\tilde{f}_n^{\lambda} \right)^2 \\ &= (2n - 1)!! \\ &:= 1 \cdot 3 \cdot 5 \cdots (2n - 1), \end{aligned}$$

the number of matchings on $1, 2, \dots, 2n$.