

A Zonotope Associated with Graphical Degree Sequences

RICHARD P. STANLEY

ABSTRACT. Let D_n denote the convex hull in \mathbf{R}^n of all (ordered) degree sequences of simple n -vertex graphs. Using the fact that D_n is a zonotope, an explicit generating function is found for the number of these degree sequences. The f -vector of D_n is found using Zaslavsky's theory of signed graph colorings. Finally we give a generalization based on a result of Fulkerson, Hoffman, and MacAndrew.

1. Introduction

Let G be a simple graph (i.e., no loops or multiple edges) with vertex set $[n] := \{1, 2, \dots, n\}$. Let $\deg(i)$ denote the degree of vertex i . We call the sequence $d(G) := (\deg(1), \dots, \deg(n))$ the *degree sequence* of G . Note that our degree sequences are *ordered*; we do not require, as is often done, that the degrees are listed in descending order. Regard the vector $d(G)$ as a point in the vector space \mathbf{R}^n , and define

$$D_n = \text{conv}\{d(G) : G \text{ is a simple graph with vertex set } [n]\},$$

where conv denotes convex hull. Thus D_n is a convex polytope, first considered by Koren [6] (suggested to him by M. Perles) and called the *polytope of degree sequences* (of length n). Koren determined the vertices of D_n and gave a system of (redundant) linear inequalities which determine D_n . Subsequently Peled and Srinivasan [8] obtained considerable further information about D_n , including a description of its facets and edges.

It follows from the Erdős-Gallai inequalities (explained in more detail below) that an integer point $(d_1, \dots, d_n) \in D_n \cap \mathbf{Z}^n$ is the degree sequence of some graph G if and only if $d_1 + \dots + d_n$ is even. Peled suggested to this writer that the number of distinct (ordered) degree sequences of n -vertex graphs might be closely approximated by half the volume of D_n , and

1980 *Mathematics Subject Classification* (1985 Revision). Primary 05C30; Secondary 52A25. Partially supported by NSF grant #DMS-8401376.

asked whether it was possible to compute or estimate the volume of D_n . It turns out that D_n is a *zonotope*, a special kind of polytope (defined below) with a well-developed theory for computing the number of integer points, the volume, the f -vector, etc. Moreover, there is a slight modification of D_n , which we denote by \tilde{D}_n , which is a zonotope combinatorially equivalent to D_n , but whose integer points correspond *exactly* to the degree sequences of graphs. Using the theory of zonotopes we are able to determine exactly the number $f(n)$ of degree sequences of n -vertex graphs, viz.,

$$\sum_{n \geq 0} f(n) \frac{x^n}{n!} = \frac{1}{2} \left[\left(1 + 2 \sum_{n \geq 1} n^n \frac{x^n}{n!} \right)^{1/2} \times \left(1 - \sum_{n \geq 1} (n-1)^{n-1} \frac{x^n}{n!} \right) + 1 \right] e^{\sum_{n \geq 1} n^{n-2} x^n / n!}.$$

Moreover, using the connection between the zonotope \tilde{D}_n and Zaslavsky's theory of signed coloring [14], we are able to compute exactly the f -vector of \tilde{D}_n (or of D_n).

2. The Ehrhart polynomial of an integer zonotope

A *zonotope* is by definition a (Minkowski) sum of closed line segments. In other words, if L_1, \dots, L_r is a set of closed line segments in \mathbf{R}^n , then the corresponding zonotope is given by

$$(1) \quad Z(L_1, \dots, L_r) = \{\alpha_1 + \dots + \alpha_r : \alpha_1 \in L_1, \dots, \alpha_r \in L_r\}.$$

If L_i joins the origin with the vector $\beta_i \in \mathbf{R}^n$, then we also write $Z(\beta_1, \dots, \beta_r)$ for $Z(L_1, \dots, L_r)$.

For any positive integer q and any convex polytope \mathcal{P} with integer vertices, let $i(\mathcal{P}, q)$ denote the number of points $\alpha \in \mathcal{P}$ satisfying $q\alpha \in \mathbf{Z}^n$. Then $i(\mathcal{P}, q)$ is a polynomial function of q of degree $\dim \mathcal{P}$, called the *Ehrhart polynomial* of \mathcal{P} . Thus $i(\mathcal{P}, 1)$ is equal to the number of integer points in \mathcal{P} . Moreover, if $\mathcal{P} \subset \mathbf{R}^n$, then the coefficient of q^n in $i(\mathcal{P}, q)$ is equal to the n -dimensional volume of \mathcal{P} . For these and other basic facts concerning Ehrhart polynomials, see, e.g., [11, pp. 235–241]. The following lemma is a refinement of a result of Shephard [9, Theorem 54]. Define a *half-open cube* C to be a Minkowski sum of linearly independent half-open line segments L_1, \dots, L_s . In other words, if $\alpha_1, \dots, \alpha_s, \beta_1, \dots, \beta_s \in \mathbf{R}^n$ and $\beta_1 - \alpha_1, \dots, \beta_s - \alpha_s$ are linearly independent, then C has the form

$$C = \{a_1\alpha_1 + (1-a_1)\beta_1 + a_2\alpha_2 + (1-a_2)\beta_2 + \dots + a_s\alpha_s + (1-a_s)\beta_s : 0 \leq a_i < 1\}.$$

We call the vectors $\gamma_i := \beta_i - \alpha_i$ the *generating vectors* of C . (If $s = 0$ then C is a single point, whose set of generating vectors is empty.)

2.1. LEMMA. *Let L_i be the line segment connecting the origin to $\gamma_i \in \mathbf{R}^n$, $1 \leq i \leq r$. Then the zonotope $Z = Z(L_1, \dots, L_r) = Z(\gamma_1, \dots, \gamma_r)$ is a disjoint union $Z = \bigsqcup_X C_X$ of half-open cubes C_X , where (a) X ranges over all linearly independent subsets of $\{\gamma_1, \dots, \gamma_r\}$, and (b) if $X = \{\delta_1, \dots, \delta_s\}$ then there are signs $e_i \in \{-1, 1\}$ such that C_X is generated by $e_1\delta_1, \dots, e_s\delta_s$. Moreover, if each $\gamma_i \in \mathbf{Z}^n$ then the vertices of each C_X lie in \mathbf{Z}^n .*

SKETCH OF PROOF (suggested by G. Ziegler). The proof is a ‘‘Tutte-Grothendieck argument’’, in the sense of [2]. We proceed by induction on r . The lemma is clear for $r = 0$, when Z consists of a single point. Assume now the case for $r - 1$. Let $Z' = Z(L_1, \dots, L_{r-1})$, so Z' has a desired decomposition $Z' = \bigsqcup_X C_X$, where X ranges over all linearly independent subsets of $\{\gamma_1, \dots, \gamma_{r-1}\}$. Now orthogonally project Z' to a hyperplane orthogonal to L_r . We obtain a zonotope \bar{Z} generated by the projections $\bar{L}_1, \dots, \bar{L}_{r-1}$ of the line segments L_1, \dots, L_{r-1} . By induction decompose $\bar{Z} = \bigsqcup_{\bar{Y}} C_{\bar{Y}}$, where the sets \bar{Y} are projections of subsets Y of $\{\gamma_1, \dots, \gamma_{r-1}\}$ for which $Y \cup \{\gamma_r\}$ is linearly independent. Now ‘‘lift’’ this decomposition back up to \mathbf{R}^n . Each half-open cube $C_{\bar{Y}}$ is lifted to a product $(O, \gamma_r] \times C_Y$, where $(O, \gamma_r]$ denotes the half-open interval from the origin O to γ_r which excludes O and includes γ_r . Then

$$Z = \left(\bigcup_X C_X \right) \cup \left(\bigcup_Y (O, \gamma_r] \times C_Y \right)$$

gives a desired decomposition of Z . \square

The next theorem is a basic result about the Ehrhart polynomial of an integer zonotope. It was stated without proof in [10, Example 3.1; 11, Exercise 4.31]. The special case of the volume of a zonotope $Z \subset \mathbf{R}^n$ (the coefficient of q^n in $i(Z, q)$) is due to McMullen (see [9, (57)]) and also appears in [7].

2.2. THEOREM. *Let $\beta_1, \dots, \beta_r \in \mathbf{Z}^n$. Let Z denote the zonotope $Z(\beta_1, \dots, \beta_r)$. Then*

$$i(Z, q) = \sum_X h(X)q^{|X|},$$

where X ranges over all linearly independent subsets of $\{\beta_1, \dots, \beta_r\}$, and where $h(X)$ denotes the greatest common divisor of all minors of size $|X|$ of the matrix whose rows are the elements of X .

PROOF. By the preceding lemma, we have $Z = \bigsqcup_X C_X$, where X runs over all linearly independent subsets $\{\delta_1, \dots, \delta_s\}$ of $\{\beta_1, \dots, \beta_r\}$, and where C_X is a half-open cube with integer vertices generated by $\pm\delta_1, \dots, \pm\delta_s$ for some choice of signs. Suppose C is any half-open cube with integer vertices

generated by vectors $\zeta_1, \dots, \zeta_s \in \mathbf{Z}^n$. Let $Y = \{\zeta_1, \dots, \zeta_s\}$. Let Λ be the (additive) abelian group of integer vectors in the subspace of \mathbf{R}^n spanned by Y , so $\Lambda \cong \mathbf{Z}^s$. Let Γ be the subgroup of Λ generated by Y . By standard properties of determinants we have $h(Y) = [\Lambda : \Gamma]$, the index of Γ in Λ . On the other hand, it is easy to see that $C \cap \Lambda$ is a set of coset representatives for Γ in Λ . Hence $i(C, 1) = \#(C \cap \Lambda) = h(Y)$. By a simple scaling argument it follows that $i(C, q) = h(Y)q^{|Y|}$ for any integer $q \geq 1$. Hence

$$i(Z, q) = \sum_X i(C_X, q) = \sum_X h(X)q^{|X|}. \quad \square$$

3. Counting degree sequences

We will now apply Theorem 2.2 to a modification \tilde{D}_n of the polytope D_n . For a graph G on the vertex set $[n]$ with degree sequence $d(G) = (d_1, \dots, d_n)$, define the *extended degree sequence*

$$\tilde{d}(G) = (d_1, \dots, d_n, \frac{1}{2}(d_1 + \dots + d_n)).$$

Then define the *polytope of extended degree sequences*,

$$\tilde{D}_n = \text{conv}\{\tilde{d}(G) : G \text{ is a simple graph on the vertex set } [n]\} \subset \mathbf{R}^{n+1}.$$

Since the last component of vectors in \tilde{D}_n is a fixed linear combination of the first n components, it follows that D_n and \tilde{D}_n are linearly equivalent and hence combinatorially equivalent. However (and this is the point of defining \tilde{D}_n), they do not have the same Ehrhart polynomial.

Let us recall the Erdős-Gallai conditions (see any graph theory text) characterizing the degree sequences of simple graphs: A vector (d_1, \dots, d_n) of positive integers d_i satisfying $d_1 \geq \dots \geq d_n$ is the degree sequence of some simple graph G if and only if the following two conditions hold:

$$(EG1) \quad \sum_{i=1}^j d_i - j(j-1) \leq \sum_{i=j+1}^n \min(j, d_i), \quad 1 \leq j \leq n.$$

$$(EG2) \quad d_1 + \dots + d_n \text{ is even.}$$

From these conditions it is easy to deduce (see [6]) that the condition for an arbitrary vector (d_1, \dots, d_n) of positive integers (i.e., not necessarily satisfying $d_1 \geq \dots \geq d_n$) to be the degree sequence of a simple graph is given by a system of linear inequalities, together with (EG2). (These conditions in fact follow from an earlier result of Tutte [12] and are also obtained in a more general context in [4, Theorem 2.1].) It follows that $(d_1, \dots, d_n) \in \mathbf{Z}^n$ is a degree sequence if and only if $(d_1, \dots, d_n) \in D_n$ and $d_1 + \dots + d_n$ is even. Hence from the definition of \tilde{D}_n we conclude:

3.1. PROPOSITION. *There is a one-to-one correspondence between degree sequences $d(G)$ of simple graphs G on the vertex set $[n]$ and integer points in \tilde{D}_n , given by $d(G) \leftrightarrow \tilde{d}(G)$.*

More generally, an analogue of the Erdős-Gallai condition for multigraphs (graphs with repeated edges, but no loops) with edge multiplicity bounded by q yields the following result (which is generalized further in Proposition 5.2).

3.2. PROPOSITION. *Let q be a positive integer. Then $i(\tilde{D}_n, q)$ is equal to the number of distinct degree sequences of multigraphs on the vertex set $[n]$ such that no edge multiplicity can exceed q .*

Now let e_i denote the i th unit coordinate vector in \mathbf{R}^n , and write $e_{ij} = e_i + e_j$. Let \tilde{e}_{ij} be the vector in \mathbf{R}^{n+1} obtained by adjoining to e_{ij} an $(n+1)$ st coordinate equal to 1. Clearly for a graph G on the vertex set $[n]$,

$$d(G) = \sum e_{ij}, \quad \tilde{d}(G) = \sum \tilde{e}_{ij},$$

where both sums range over all edges $\{i, j\}$ of G . From this it follows easily that D_n and \tilde{D}_n are zonotopes, given explicitly by

$$(2) \quad \begin{aligned} D_n &= Z(e_{ij} : 1 \leq i < j \leq n), \\ \tilde{D}_n &= Z(\tilde{e}_{ij} : 1 \leq i < j \leq n). \end{aligned}$$

(Of course, since D_n and \tilde{D}_n are linearly equivalent, it follows that one is a zonotope if and only if the other is.)

Now define a *quasitree* to be a connected graph which is either a tree or has exactly one cycle, which is of odd length. A *quasiforest* is a graph whose components are all quasitrees. We have come to the main result of this section.

3.3. THEOREM. *The Ehrhart polynomial $i(\tilde{D}_n, q)$ of \tilde{D}_n is given by*

$$i(\tilde{D}_n, q) = a_n(n)q^n + a_n(n-1)q^{n-1} + \dots + a_n(0),$$

where

$$a_n(i) = \sum_X \max\{1, 2^{c(X)-1}\};$$

here X ranges over all quasiforests with i edges on the vertex set $[n]$, and $c(X)$ denotes the number of (odd) cycles of X .

PROOF. By Theorem 2.2 and (2), we have

$$a_n(i) = \sum_X h(X),$$

where X ranges over all i -element linearly independent subsets of $\tilde{E}_n := \{\tilde{e}_{ij} : 1 \leq i < j \leq n\}$, and $h(X)$ has the meaning of Theorem 2.2. Let us identify a subset X of \tilde{E}_n with the graph on the vertex set $[n]$ which has an edge $\{i, j\}$ if and only if $\tilde{e}_{ij} \in X$. When is X linearly independent? We could appeal to Zaslavsky's extensive theory of signed graphs [13] to

answer this question, but it is also easy to proceed directly. If X is linearly independent, then every component Y of X with i vertices may contain at most i edges, since any $i + 1$ vectors in \mathbf{R}^i are dependent. Thus every component Y of X is either a tree or contains one cycle. It is easily checked that a connected graph Y with at most one cycle is linearly independent if and only if Y is a tree or the unique cycle of Y has odd length. Thus we have that X is a quasiforest.

Assume then that X is a quasiforest, and set $m = |X|$. We need to compute $h(X)$, the gcd of the $m \times m$ minors of the matrix M whose rows are the vectors $\tilde{e}_{ij} \in X$. Thus the rows of M are indexed by the edges of X and the columns by the vertices, together with a last column of 1s. An $m \times m$ minor of M corresponds to choosing either m vertices of X (i.e., m numbers from $[n]$) or $m - 1$ vertices of X and a final column of 1s. In the first case, in order for the columns to be linearly independent we must choose all vertices from any component of X with an odd cycle, and all but one vertex from any component of X which is a tree. Each odd cycle contributes to a factor ± 2 to the minor, and we obtain a value $\pm 2^{c(X)}$.

Now consider the submatrix N obtained by choosing $m - 1$ vertices of X and a final column of 1s. Again, in order for the columns to be linearly independent we must choose our $m - 1$ vertices in one of the following ways: (a) choose all but one vertex from every component Y which is a tree, choose all but one vertex from some component T_0 which has an odd cycle, and choose every vertex from the remaining components T with one odd cycle, or (b) choose all but two adjacent vertices i and j from some component which is a tree, choose all but one vertex from the remaining components which are trees, and choose every vertex from the components with one odd cycle.

In case (a), in the submatrix N subtract one-half of every column except the last from the last column. Every entry of the last column is now 0 or $1/2$. Factor out $1/2$ from the last column. The resulting matrix N' is the incidence matrix of a graph X' for which every component has exactly one cycle, which is of odd length. Moreover, X' has the same number of odd cycles as X . Hence $\det N' = \pm 2^{c(X)}$, so $\det N = \pm 2^{c(X)-1}$.

In case (b), there is a row which is all 0s except a 1 in the last column. Hence we may delete this row and the last column without affecting the value of the minor (except possibly for sign). Let X' denote the graph X with edge $\{i, j\}$ deleted (but retain all vertices). We now have a matrix obtained from the incidence matrix of X' by deleting a column (vertex) from every component which is a tree. Since X' still has $c(X)$ odd cycles, it follows that the value of the minor is $\pm 2^{c(X)}$.

Thus we have seen that all nonzero minors of M are equal to $\pm 2^{c(X)}$ or $\pm 2^{c(X)-1}$. The latter value is possible if and only if $c(X) \geq 1$. Hence the gcd of the minors of M is given by $\max\{1, 2^{c(X)-1}\}$, and the proof follows. \square

3.4. COROLLARY. *The number $f(n)$ of distinct degree sequences of simple graphs on the vertex set $[n]$ is given by*

$$(3) \quad f(n) = \sum_X \max\{1, 2^{c(X)-1}\},$$

where X ranges over all quasiforests on the vertex set $[n]$, and $c(X)$ denotes the number of (odd) cycles of X .

PROOF. Put $q = 1$ in Theorem 3.3. \square

It would be interesting to find a direct combinatorial proof of (3). An analogous problem to determining $f(n)$ is to find the number $g(n)$ of distinct outdegree sequences of orientations of the complete graph K_n . Using the theory of zonotopes as we have done here, it was shown in [10, Example 3.1] (see also [11, Exercise 4.32]) that $g(n)$ is equal to the number of (labelled) forests on n vertices. A combinatorial proof of this result was subsequently given by Kleitman and Winston [5]. Perhaps their techniques can be used to establish (3). However, while the enumeration of outdegree sequences can be extended to orientations of any undirected graph (as discussed in the previous three references), an analogous generalization for degree sequences of subgraphs of a given graph G seems only to hold for a special class of graphs G (see §5 for further details).

It is a fairly routine exercise in enumerative combinatorics to convert Theorem 3.3 into a generating function for $i(\tilde{D}_n, q)$.

3.5. PROPOSITION. *We have*

$$\sum_{n \geq 0} i(\tilde{D}_n, q) \frac{x^n}{n!} = \frac{1}{2} \left[\left(1 + 2 \sum_{n \geq 1} n^n \frac{q^n x^n}{n!} \right)^{1/2} \times \left(1 - \sum_{n \geq 1} (n-1)^{n-1} \frac{q^n x^n}{n!} \right) + 1 \right] e^{\sum_{n \geq 1} n^{n-2} q^{n-1} x^n / n!}.$$

Here and below we set $0^0 = 1$ (which arises when $n = 1$ in the second sum on the right-hand side).

PROOF. We assume knowledge of the theory of exponential generating functions, in particular the exponential formula, as expounded for instance in [3, Theorem 4.3]. The number of rooted trees on n vertices is n^{n-1} , with exponential generating function

$$R(x) = \sum_{n \geq 1} n^{n-1} \frac{x^n}{n!}.$$

Hence the exponential generating function for k -tuples of rooted trees is $R(x)^k$ and so for undirected k -cycles of rooted trees (i.e., graphs with exactly one cycle, which is of length $k \geq 3$) is $R(x)^k / 2k$.

Let $h(j, n)$ be the number of graphs on the vertex set $[n]$ such that every component has exactly one cycle, which is of odd length ≥ 3 , and with j

cycles. (Such graphs have exactly n edges.) Then by the exponential formula we have

$$\begin{aligned} \sum_{j, n \geq 0} h(j, n) \frac{t^j x^n}{n!} &= \exp \sum_{k \geq 1} \frac{t}{2(2k+1)} R(x)^{2k+1} \\ &= \exp \frac{t}{2} \left[\frac{1}{2} (\log(1 - R(x))^{-1} - \log(1 + R(x))^{-1}) - R(x) \right] \\ &= \left(\frac{1 + R(x)}{1 - R(x)} \right)^{t/4} e^{-tR(x)/2}. \end{aligned}$$

Thus,

$$\begin{aligned} 1 + \sum_{j, n \geq 1} 2^{j-1} h(j, n) \frac{x^n}{n!} &= 1 + \frac{1}{2} \left[\left(\frac{1 + R(x)}{1 - R(x)} \right)^{1/2} e^{-R(x)} - 1 \right] \\ &= \frac{1}{2} \left[\left(-1 + \frac{2}{1 - R(x)} \right)^{1/2} e^{-R(x)} + 1 \right]. \end{aligned}$$

It is well known (and easily deduced from $R(x) = xe^{R(x)}$) that

$$\frac{1}{1 - R(x)} = \sum_{n \geq 0} n^n \frac{x^n}{n!}, \quad e^{-R(x)} = 1 - \sum_{n \geq 1} (n-1)^{n-1} \frac{x^n}{n!},$$

so we get

$$\begin{aligned} (4) \quad &1 + \sum_{j, n \geq 1} 2^{j-1} h(j, n) \frac{x^n}{n!} \\ &= \frac{1}{2} \left[\left(1 + 2 \sum_{n \geq 1} n^n \frac{x^n}{n!} \right)^{1/2} \left(1 - \sum_{n \geq 1} (n-1)^{n-1} \frac{x^n}{n!} \right) + 1 \right]. \end{aligned}$$

There are n^{n-2} free (i.e., unrooted) trees with n labelled vertices, each with $n-1$ edges. Hence if $r(i, n)$ denotes the number of forests with i edges on the vertex set $[n]$, then

$$\sum_{i, n \geq 0} r(i, n) \frac{q^i x^n}{n!} = \exp \sum_{n \geq 1} n^{n-2} \frac{q^{n-1} x^n}{n!}.$$

Since all graphs enumerated by $h(n, j)$ have exactly n edges, there follows

$$\sum_{n \geq 0} i(\tilde{D}_n, q) \frac{x^n}{n!} = \left[1 + \sum_{j, n \geq 1} 2^{j-1} h(j, n) \frac{(qx)^n}{n!} \right] \cdot \exp \sum_{n \geq 1} n^{n-2} \frac{q^{n-1} x^n}{n!}.$$

Comparing with (4) completes the proof. \square

Putting $q = 1$ in Proposition 3.5 yields:

3.6. COROLLARY. *We have*

$$\sum_{n \geq 0} f(n) \frac{x^n}{n!} = \frac{1}{2} \left[\left(1 + 2 \sum_{n \geq 1} n \frac{x^n}{n!} \right)^{1/2} \times \left(1 - \sum_{n \geq 1} (n-1) \frac{x^n}{n!} \right) + 1 \right] e^{\sum_{n \geq 1} n^{n-2} x^n / n!}.$$

Some small values of $i_n := i(\tilde{D}_n, q)$ are given by:

$$i_1 = 1$$

$$i_2 = 1 + q$$

$$i_3 = 1 + 3q + 3q^2 + q^3$$

$$i_4 = 1 + 6q + 15q^2 + 20q^3 + 12q^4$$

$$i_5 = 1 + 10q + 45q^2 + 120q^3 + 195q^4 + 162q^5$$

$$i_6 = 1 + 15q + 105q^2 + 455q^3 + 1320q^4 + 2508q^5 + 2540q^6$$

$$i_7 = 1 + 21q + 210q^2 + 1330q^3 + 5880q^4 + 18564q^5 + 39809q^6 + 46035q^7$$

$$i_8 = 1 + 28q + 378q^2 + 3276q^3 + 20265q^4 + 93240q^5 + 317800q^6 + 749200q^7 + 951552q^8.$$

Moreover,

$$f(1) = 1$$

$$f(2) = 2$$

$$f(3) = 8$$

$$f(4) = 54$$

$$f(5) = 533$$

$$f(6) = 6944$$

$$f(7) = 111850$$

$$f(8) = 2135740$$

$$f(9) = 47003045$$

$$f(10) = 1168832808.$$

The coefficient of q^n in $i(\tilde{D}_n, q)$ is the n -dimensional *relative volume* $V(\tilde{D}_n)$ of \tilde{D}_n (see, e.g., [10, p. 335] or [11, pp. 238–239] for the definition). Putting $1/q$ for q and qx for x in Proposition 3.5 and then setting $q = 0$ yields:

3.7. COROLLARY. *We have*

$$\sum_{n \geq 0} V(\tilde{D}_n) \frac{x^n}{n!} = \frac{1}{2} \left[\left(1 + 2 \sum_{n \geq 1} n^n \frac{x^n}{n!} \right)^{1/2} \left(1 - \sum_{n \geq 1} (n-1)^{n-1} \frac{x^n}{n!} \right) + 1 \right].$$

We also have that

$$(5) \quad V(\tilde{D}_n) = \sum_X 2^{c(X)-1},$$

where X ranges over all graphs on the vertex set $[n]$ for which every component has exactly one cycle, which is of odd length, and where $c(X)$ denotes the number of (odd) cycles of X . Moreover, in regard to Peled's question about the volume of D_n , we have by similar reasoning that

$$V(D_n) = \sum_X 2^{c(X)},$$

summed over the same range as (5), so $V(D_n) = 2V(\tilde{D}_n)$. Moreover, $V(D_n)$ is the ordinary volume of D_n for $n \geq 3$.

4. The f -vector of D_n and \tilde{D}_n

We mentioned in §1 that Koren, Peled, and Srinivasan obtained a description of the vertices, edges, and facets of D_n (or \tilde{D}_n). Beissinger and Peled [1] used the description of the vertices to obtain a formula for the number of vertices (see equation (7)). Here we extend this result to a determination of the entire f -vector $f(D_n) = (f_0(D_n), \dots, f_n(D_n)) = (f_0, \dots, f_n)$, where D_n has f_i i -dimensional faces. (Normally the f -vector of a d -polytope only includes f_i for $0 \leq i \leq d-1$, but it will be convenient for us to allow $0 \leq i \leq n$.) Thus, $f(D_1) = (1, 0)$, $f(D_2) = (2, 1, 0)$, and $f(D_3) = (8, 12, 6, 1)$, the latter since D_3 is a 3-cube.

Our derivation will be based on a result of Zaslavsky concerning the coloring of signed graphs. Let $-K_n$ denote the *negative complete signed graph*, i.e., the complete graph K_n with every edge labelled with a minus sign [13, §7D]. The matroid $M = M(-K_n)$ which Zaslavsky associates with $-K_n$ is just the linear matroid $\{e_{ij} : 1 \leq i < j \leq n\} \subset \mathbf{R}^n$. Hence by (2), Zaslavsky's zonotope $Z[-K_n]$ (defined in [14, p. 226]) is just D_n .

In [14, pp. 217–218], Zaslavsky defines a *signed coloring* of $-K_n$ (more generally, Zaslavsky deals with an arbitrary signed graph Σ) in $\mu \geq 0$ colors to be a function $k : N \rightarrow [-\mu, \mu] = \{-\mu, -\mu+1, \dots, 0, \dots, \mu-1, \mu\}$, where $N = [n]$ denotes the vertex set of $-K_n$. The *set of impropriety* $I(k)$ of k consists of all edges $\{i, j\}$ of $-K_n$ for which $k(i) = -k(j)$ [14, p. 218]. Call such edges *improper*. The *rank* of $I(k)$ is defined by $\text{rk} I(k) = n - b(I(k))$, where $b(I(k))$ is the number of bipartite components of $I(k)$ (considered as a graph on the vertex set $[n]$) [14, p. 216]. The *Whitney*

polynomial of $-K_n$ is defined by

$$w_{-K_n}(x, 2\mu + 1) = \sum_k x^{\text{rk } I(k)},$$

summed over all signed colorings in μ colors.

We can now state the result of Zaslavsky [14, Corollary 4.1''] (specialized to $\Sigma = -K_n$ and $Z[\Sigma] = D_n$), which will be our main tool for evaluating the f -vector $f(D_n)$.

4.1. PROPOSITION. *We have*

$$\sum_{i=0}^n f_i(D_n)x^i = (-1)^n w_{-K_n}(-x, -1).$$

The main result of this section is the following:

4.2. THEOREM. *We have*

$$\sum_{n \geq 0} \sum_{i=0}^n f_i(D_n) \frac{x^i t^n}{n!} = \frac{e^{xt} - (1+x)t - x(1+x)\frac{t^2}{2}}{1 + 2(e^{-t} - 1) + \frac{2}{x}(e^{xt} - 1) - \frac{1}{x}(e^{2xt} - 1)}.$$

PROOF. We may choose a signed coloring $k : N \rightarrow [-t, t]$ of $-K_n$ as follows. First choose the sets $A_i = k^{-1}\{-i, i\}$. Thus (A_0, A_1, \dots, A_μ) is a weak ordered partition of $[n]$ into $\mu + 1$ blocks, i.e., $\bigcup A_i = [n]$ and $A_i \cap A_j = \emptyset$ if $i \neq j$. ("Weak" means that we allow $A_i = \emptyset$.) Let $a_i = |A_i|$. Now for each block A_i with $1 \leq i \leq \mu$, choose a subset $B_i = k^{-1}(i)$. For two of these choices (when $A_i \neq \emptyset$), namely $B_i = A_i$ and $B_i = \emptyset$, there will be no improper edges incident to vertices in A_i , so A_i forms a_i bipartite components of $I(k)$, each consisting of a single vertex. For the remaining $2^{a_i} - 2$ choices of B_i (when $a_i \geq 1$), all edges from B_i to $A_i - B_i$ are improper; these edges form a single bipartite component of $I(k)$. There remains the case of A_0 . All edges between vertices in A_0 are improper, so we get a component of $I(k)$ isomorphic to the complete graph K_{a_0} . This component will be bipartite if and only if $a_0 = 0, 1$, or 2 . It follows that

$$(6) \quad w_{-K_n}(x, 2\mu + 1) = x^n \sum_{(A_0, \dots, A_\mu)} v(a_0) \prod_{i=1}^{\mu} (2x^{a_i} + (2^{a_i} - 2)\chi(a_i \neq 0)x^{-1}),$$

where: (a) the sum ranges over all weak ordered partitions of $[n]$ into $\mu + 1$ blocks, (b) $\chi(a_i \neq 0) = 1$ if $a_i \neq 0$ and $= 0$ if $a_i = 0$, and (c) $v(a_0)$ is defined by

$$v(a) = \begin{cases} 1 & \text{if } a = 0 \text{ or } a \geq 3, \\ x^{-1} & \text{if } a = 1 \text{ or } a = 2. \end{cases}$$

Now suppose g_0, \dots, g_μ are any functions defined on \mathbb{N} , and define (using the notation of (6))

$$h(n) = \sum_{(A_0, \dots, A_\mu)} g_0(a_0) \cdots g_\mu(a_\mu).$$

It then follows from standard properties of exponential generating functions that

$$h(n) = \left[\frac{t^n}{n!} \right] \prod_{i=0}^{\mu} \left(\sum_{j \geq 0} g_i(j) \frac{t^j}{j!} \right),$$

where $\left[\frac{t^n}{n!} \right] F(t)$ denotes the coefficient of $t^n/n!$ in the generating function $F(t)$. Hence from (6) we have

$$\begin{aligned} W_{K_n}(x, 2\mu + 1) &= x^n \left[\frac{t^n}{n!} \right] \left(x^{-1}t + x^{-1}\frac{t^2}{2} + \sum_{j \geq 3} \frac{t^j}{j!} \right) \left(1 + \sum_{j \geq 1} (2x^{-j} + (2^j - 2)x^{-1}) \frac{t^j}{j!} \right)^\mu \\ &= \left[\frac{t^n}{n!} \right] \left(1 + \frac{xt^2}{2} + \sum_{j \geq 3} \frac{(xt)^j}{j!} \right) \left(1 + \sum_{j \geq 1} (2x^{-j} + (2^j - 2)x^{-1}) \frac{(xj)^j}{j!} \right)^\mu \\ &= \left[\frac{t^n}{n!} \right] \left((1-x)t + x(1-x)\frac{t^2}{2} + e^{xt} \right) \\ &\quad \times \left(1 + 2(e^t - 1) - \frac{2}{x}(e^{xt} - 1) + \frac{1}{x}(e^{2xt} - 1) \right)^\mu. \end{aligned}$$

Substituting $-x$ for x , $-t$ for t , and -1 for μ yields

$$\begin{aligned} (-1)^n w_{-K_n}(-x, -1) &= \left[\frac{t^n}{n!} \right] \left(e^{xt} - (1+x)t - x(1+x)\frac{t^2}{2} \right) \\ &\quad \times \left(1 + 2(e^{-t} - 1) + \frac{2}{x}(e^{xt} - 1) - \frac{1}{x}(e^{2xt} - 1) \right)^{-1}. \end{aligned}$$

The proof follows from Proposition 4.1. \square

From Theorem 4.2 one can compute:

$$f(D_1) = (1, 0)$$

$$f(D_2) = (2, 1, 0)$$

$$f(D_3) = (8, 12, 6, 1)$$

$$f(D_4) = (46, 108, 84, 22, 1)$$

$$f(D_5) = (332, 1020, 1080, 450, 60, 1)$$

$$f(D_6) = (2874, 10830, 14880, 9160, 2460, 224, 1)$$

$$f(D_7) = (29874, 129486, 220920, 182770, 75670, 14238, 882, 1)$$

$$f(D_8) = (334982, 1726648, 3529344, 3679872, 2074660, 610288, 81144, 3322, 1).$$

Putting $x = 0$ in Theorem 4.2 yields

$$(7) \quad \sum_{n \geq 0} f_0(D_n) \frac{t^n}{n!} = \frac{1-t}{-1+2e^{-t}} = \frac{(1-t)e^t}{2-e^t},$$

agreeing with a result of Beissinger and Peled [1, p. 216] mentioned at the beginning of this section.

5. A generalization

We wish to generalize Theorem 3.3 by considering degree sequences of graphs which are contained in a fixed graph G . More precisely, let G be a multigraph (i.e., allowing multiple edges, but not loops) on the vertex set $[n]$. Define a convex polytope $\tilde{D}(G) \subset \mathbf{R}^{n+1}$ by

$$\tilde{D}(G) = \text{conv}\{\tilde{d}(H) : H \text{ is a spanning subgraph of } G\}.$$

Thus $\tilde{D}_n = \tilde{D}(K_n)$, where K_n denotes the complete graph on $[n]$. It is easily seen that

$$\tilde{D}_n = Z(\tilde{e}_{ij} : \{i, j\} \text{ is an edge of } G).$$

(Here we take \tilde{e}_{ij} a total of m times if there are m edges between i and j .) The following generalization of Theorem 3.3 is proved in exactly the same way as Theorem 3.3.

5.1. THEOREM. *The Ehrhart polynomial $i(\tilde{D}(G), q)$ of $\tilde{D}(G)$ is given by*

$$i(\tilde{D}(G), q) = a_G(n)q^n + a_G(n-1)q^{n-1} + \dots + a_G(0),$$

where

$$a_G(i) = \sum_X \max\{1, 2^{c(X)-1}\};$$

here X ranges over all spanning quasiforests of G with i edges, and $c(X)$ denotes the number of (odd) cycles of X .

We would like to interpret $i(\tilde{D}(G), q)$ in terms of degree sequences analogously to Proposition 3.2. We therefore need to know when Proposition 3.1 extends to $\tilde{D}(G)$. Call G an *FHM-graph* (named after the authors of [4]) if the following property holds:

(FHM). No induced subgraph of G consists of two vertex-disjoint odd cycles (with no other edges). (Equivalently, every induced subgraph of G has at most one nonbipartite component.)

Given a multigraph G on the vertex set $[n]$, let

$$\Delta(G) = \{d(H) : H \text{ is a spanning subgraph of } G\}.$$

Define a map $\varphi_G : \Delta(G) \rightarrow \mathbf{Z}^{n+1}$ by $\varphi_G(d(H)) = \tilde{d}(H)$. Clearly φ_G is one-to-one. It is immediate from the definition of $\tilde{D}(G)$ that

$$(8) \quad \varphi_G(\Delta(G)) \subseteq \tilde{D}(G) \cap \mathbf{Z}^{n+1}.$$

5.2. PROPOSITION. *We have $\varphi_G(\Delta(G)) = \tilde{D}(G) \cap \mathbf{Z}^{n+1}$ if and only if G is an FHM-graph.*

PROOF. The “if” part follows from [4, Theorem 2.1], while the “only if” part (which is easy) follows from the last paragraph on p. 170 of [4]. \square^1

¹I am grateful to L. Lovász for bringing the paper [4] to my attention.

Now given any multigraph G on $[n]$ and any integer $q \geq 1$, define $G^{(q)}$ to be the multigraph obtained from G by replacing each edge by q edges, and let $f(G^{(q)})$ denote the number of distinct degree sequences of spanning subgraphs of $G^{(q)}$.

5.3. THEOREM. *For any multigraph G on $[n]$ and any integer $q \geq 1$, we have*

$$f(G^{(q)}) \leq i(\tilde{D}(G), q).$$

Equality holds if and only if G is an FHM-graph.

PROOF. From the definition of FHM-graph we see that for any integer $q \geq 1$, G is an FHM-graph if and only if $G^{(q)}$ is. Moreover, by the definition of $\tilde{D}(G)$ we have $\tilde{D}(G^{(q)}) = q\tilde{D}(G)$, where for any polytope \mathcal{P} we define

$$q\mathcal{P} = \{q\alpha : \alpha \in \mathcal{P}\}.$$

Since

$$i(\tilde{D}(G), q) = \#(q\tilde{D}(G) \cap \mathbf{Z}^{n+1}) = \#(\tilde{D}(G^{(q)}) \cap \mathbf{Z}^{n+1}),$$

the proof follows from (8) and Proposition 5.2 (and the fact that $\varphi_{G^{(q)}}$ is one-to-one). \square

A special case of FHM-graphs is bipartite graphs. Theorems 5.1 and 5.3 yield for these graphs the following result.

5.4. COROLLARY. *Let G be a bipartite (multi)graph on the vertex set $[n]$. Then*

$$f(G^{(q)}) = b_G(n-1)q^{n-1} + b_G(n-2)q^{n-2} + \cdots + b_G(0),$$

where $b_G(i)$ is the number of spanning forests of G with i edges. In particular, $f(G)$ is equal to the number of spanning forests of G .

We mentioned after Corollary 3.4 that the number $g(G)$ of outdegree sequences of orientations of any graph (or multigraph) G is equal to the number of spanning forests of G , and that this result has a combinatorial proof. Hence by the preceding corollary we have $f(G) = g(G)$ when G is bipartite. It is easy to give a combinatorial proof of this fact. Namely, if G has vertex bipartition (A, B) and if D is an orientation of G , then let $\sigma(D)$ be the spanning subgraph of G consisting of those edges which are oriented from A to B in D . Then D and D' have the same outdegree sequence if and only if $\sigma(D)$ and $\sigma(D')$ have the same degree sequence, yielding the desired bijection. Hence the combinatorial proof [5] yields a combinatorial proof of Corollary 5.4 when $q = 1$, and this leads easily to a combinatorial proof for any q .

Combining Theorems 5.1 and 5.3 when $q = 1$ yields that for an FHM-graph G ,

$$(9) \quad f(G) = \sum_X \max\{1, 2^{c(X)-1}\},$$

where X ranges over all spanning quasiforests of G . It would be interesting to find a generalization of (9) valid for all graphs G . While we have been unable to do this, there is a more general class of graphs than FHM-graphs which we can handle. Recall that a *closed trail* T of length m in a graph G is a sequence $v_0 e_1 v_1 e_2 v_2 \cdots e_m v_m$ such that each v_i is a vertex, each e_i is an edge, $v_0 = v_m$, any two adjacent terms are incident in G , and finally all the e_i 's are distinct. Call the closed trail T *even* if m is even.

5.5. PROPOSITION. *Let e be an edge of the graph G . Then $f(G) = 2f(G - e)$ if and only if e does not belong to an even closed trail of G .*

PROOF. Note that $f(G) = 2f(G - e)$ if and only if there do not exist spanning subgraphs H, H' of G such that $d(H) = d(H')$, e is an edge of H , and e is not an edge of H' . Suppose e is an edge of the even closed trail $v_0 e_1 v_1 \cdots e_{2r} v_{2r}$. Let H have edges $e_1, e_3, \dots, e_{2r-1}$, and H' have edges e_2, e_4, \dots, e_{2r} . Then $d(H) = d(H')$, and e is an edge of exactly one of H and H' . Thus $f(G) \neq 2f(G - e)$.

Conversely, suppose H and H' are spanning subgraphs of G such that $d(H) = d(H')$, and let e be an edge of H but not of H' . Let B be the set of edges of H which are not edges of H' , and let R be the set of edges of H' which are not edges of H . Regard the edges in B as "blue" and in R as "red". Consider the graph G' whose edges are $B \cup R$. Every vertex G' is incident to the same number of blue edges as red edges. It follows easily that there is a closed trail T containing e whose edge alternate blue and red. Hence T is even, as desired. \square

The point of Proposition 5.5 is that even though G may not be an FHM-graph, after we remove all edges not belonging to even closed trails the resulting graph may then be an FHM-graph. Thus equation (9) and Proposition 5.5 gives us a formula for $f(G)$. We also may assume G is connected, since clearly $f(G + H) = f(G)f(H)$, where $G + H$ denotes the disjoint union of G and H .

REFERENCES

1. J. S. Beissinger and U. N. Peled, *Enumeration of labelled threshold graphs and a theorem of Frobenius involving Eulerian polynomials*, Graphs Combin. **3** (1987), 213–219.
2. T. Brylawski, *A decomposition for combinatorial geometries*, Trans. Amer. Math. Soc. **171** (1972), 235–282.
3. D. Foata, *La série génératrice exponentielle dans les problèmes d'énumération*, Sém. Math. Sup., Presses Univ. Montréal, 1974.
4. D. R. Fulkerson, A. J. Hoffman, and M. H. McAndrew, *Some properties of graphs with multiple edges*, Canad. J. Math. **17** (1965), 166–177.
5. D. Kleitman and K. Winston, *Forests and score vectors*, Combinatorica **1** (1981), 49–54.
6. M. Koren, *Extreme degree sequences of simple graphs*, J. Combin. Theory Ser. B **15** (1973), 213–224.
7. H. L. Montgomery and J. D. Vaaler, E3301, American Math. Monthly **96** (1989), 54.
8. U. N. Peled and M. K. Srinivasan, *The polytope of degree sequences*, Linear Algebra Appl. **114/115** (1989), 349–377.

9. G. C. Shephard, *Combinatorial properties of associated zonotopes*, *Canad. J. Math.* **26** (1974), 302–321.
10. R. Stanley, *Decompositions of rational convex polytopes*, *Ann. Discrete Math.* **6** (1980), 333–342.
11. ———, *Enumerative combinatorics*, vol. 1, Wadsworth & Brooks/Cole, Pacific Grove, Cal., 1986.
12. W. T. Tutte, *The factors of graphs*, *Canad. J. Math.* **4** (1952), 314–329.
13. T. Zaslavsky, *Signed graphs*, *Discrete Appl. Math.* **4** (1982), 47–74.
14. ———, *Signed graph coloring*, *Discrete Math.* **39** (1982), 215–228.

DEPARTMENT OF MATHEMATICS, MASSACHUSETTS INSTITUTE OF TECHNOLOGY, CAMBRIDGE,
MASSACHUSETTS 02139

E-mail address: rstan@math.mit.edu