

Your name is: \_\_\_\_\_

Grading 1  
2  
3

Total \_\_\_\_\_

Thank you for staying with 18.085. I appreciate it very much. Have a great holiday.

1) (30 pts.) Suppose we know a function  $f(x)$  between  $x = -\pi$  and  $x = \pi$ . For Fourier series we extend it *periodically* for  $|x| > \pi$ . For Fourier integrals I will extend it *by zero* for  $|x| > \pi$ .

(a) How are the Fourier series coefficients  $c_k$  related to the Fourier integral transform  $\hat{f}(k)$ ? Look at the formulas.

(b) How is this integral related to this sum?

$$\int_{-\infty}^{\infty} \hat{f}(k) e^{ikx} dk \qquad \sum_{k=-\infty}^{\infty} c_k e^{ikx}$$

It seems surprising that the integral uses all  $k$  and the sum uses only integer  $k$ . (This is the key to Shannon's Sampling Theorem! It works because  $f(x)$  is entirely given within the interval  $[-\pi, \pi]$ )

(c) If  $f(x)$  is a hat function (piecewise linear and continuous) how quickly do the coefficients  $c_k$  approach zero? What is the decay rate if  $f(x) =$  hat function *squared*?



2) (30 pts.) Suppose a function  $f(x)$  has a known Fourier integral transform  $\widehat{f}(k)$ . As usual  $-\infty < x < \infty$  and  $-\infty < k < \infty$ . Let  $B(x)$  be the box function, equal to 1 for  $|x| \leq \frac{1}{2}$  (and  $B(x) = 0$  for  $|x| > \frac{1}{2}$ ).

- (a) If you convolve  $f(x)$  with the box function  $B(x)$ , what is the Fourier integral transform  $\widehat{h}(k)$  of  $h(x) = f(x) * B(x)$ ? Do you expect  $h(x)$  to be more smooth or less smooth than  $f(x)$  (and by how much and why)?
- (b) If you consider the derivative  $h'(x) = dh/dx$ , what is its Fourier integral transform  $\widehat{h}'(k)$ ? Use the standard rule for transform of a derivative.
- (c) The convolution of  $f(x)$  with the box  $B(x)$  is actually given by

$$h(x) = \int_{-\frac{1}{2}}^{\frac{1}{2}} f(x-y)dy.$$

Go ahead and take the  $x$ -derivative using ordinary calculus (and then integrate  $f'(x-y)$  to write  $h'(x)$  without any integral). Then take the Fourier transform of this  $h'(x)$  by the usual integral formula and hopefully recover the answer in part (b).



- 3) (40 pts.) Suppose you have  $N = 8$  values  $c_0, c_1, \dots, c_7$ . Let  $c_{\text{even}}$  and  $c_{\text{odd}}$  be the “even and odd” subvectors of length 4:

$$c_{\text{even}} = c_0, c_2, c_4, c_6 \quad c_{\text{odd}} = c_1, c_3, c_5, c_7$$

- (a) With  $N = 8$  what is the complex number  $w_8 = e^{2\pi i/8}$  in its  $a + ib$  form? How is  $w_8$  related to the number  $w_4 = e^{2\pi i/4}$ ?
- (b) I am looking for a connection between the 8-point DFT of  $c$  and the 4-point DFT's of  $c_{\text{even}}$  and  $c_{\text{odd}}$ . Look at these particular components:

$$Y_1 = c_0 + c_2 w_4 + c_4 w_4^2 + c_6 w_4^3$$

$$Z_1 = c_1 + c_3 w_4 + c_5 w_4^2 + c_7 w_4^3$$

$$y_1 = c_0 + c_1 w_8 + c_2 w_8^2 + c_3 w_8^3 + c_4 w_8^4 + c_5 w_8^5 + c_6 w_8^6 + c_7 w_8^7$$

What combination of  $Y_1$  and  $Z_1$  gives  $y_1$ ? (This is the key to the Fast Fourier Transform, but just answer directly!).

- (c) Suppose  $y(n)$  is the average of the four values  $x(n), x(n - 1), x(n - 2), x(n - 3)$ :

$$y(n) = \frac{1}{4} \sum_{k=0}^3 x(n - k), \quad -\infty < n < \infty.$$

This  $y$  is a (non-cyclic) convolution of the vector  $x$  with what vector  $h$ ?

- (d) In part (c) find the relation between the functions

$$Y(\omega) = \sum y(n) e^{-in\omega} \quad \text{and} \quad X(\omega) = \sum x(n) e^{-in\omega}$$

Find an input other than  $x = (\dots, 1, -1, 1, -1, \dots)$  that leads to the output  $y = \text{all zeros}$ .

