

MODULAR DATA

PAVEL ETINGOF

The goal of this note (which contains no new results) is to give an introduction to the basic theory of modular data. We loosely follow [G] (see also [L]).¹

I am grateful to V. Drinfeld and V. Ostrik for useful explanations.

1. THE DEFINITION OF A MODULAR DATUM

1.1. Fusion rings and categories. Recall that a \mathbb{Z}_+ -ring is a ring R , finitely generated and free as a \mathbb{Z} -module, which is equipped with a distinguished basis B containing 1 in which the structure constants are nonnegative. Given a \mathbb{Z}_+ -ring R and an element $a \in R$, let $\eta(a)$ be the coefficient of 1 in the decomposition of a with respect to the distinguished basis. Recall that R is called a based ring (or a fusion ring) if there is an involution $*$: $B \rightarrow B$ (called dualization) such that $\eta(ab^*) = \delta_{ab}$ for any two elements $a, b \in B$. Clearly, if such an involution exists, it is unique.

The Grothendieck ring $R = K_0(\mathcal{C})$ of a fusion category \mathcal{C} is a fusion ring. In this situation, \mathcal{C} is said to be a categorification of R .

1.2. Modular data. Let k be an algebraically closed field of characteristic zero.

Definition 1.1. A modular datum M over k is a finite set I with a distinguished element $0 \in I$ and functions $S : I \times I \rightarrow k$ (called the S -matrix) and $T : I \rightarrow k$ (called the twist function), satisfying the following conditions:

- (1) S is symmetric, $d_i := S_{0i} \neq 0 \forall i \in I$, and $s_{00} = 1$;
- (2) The functions $X_i : I \rightarrow k$ defined by $X_i(j) := \frac{S_{ij}}{S_{0j}}$ span inside $\text{Fun}(I, k)$ a based ring R_M (in which they are the distinguished basis);
- (3) $d_i = d_{i^*}$;
- (4) If we regard S and T as linear operators on $\text{Fun}(I, k)$ (by setting $(Sf)(i) = \sum S_{ij}f(j)$, $(Tf)(i) = T(i)f(i)$), then they define a projective representation of the modular group $SL_2(\mathbb{Z})$. That is, there exists $\tau \in k^\times$ such that $(ST)^3 = \tau S^2$, and this operator commutes with T . Also, $T(0) = 1$.

The numbers d_i are called the quantum dimensions of M .

Proposition 1.2. *If \mathcal{C} is a modular tensor category, then its set of simple objects I , the S -matrix S and the twist function T form a modular datum.*

Definition 1.3. A categorification of a modular datum M is a categorification \mathcal{C} of the based ring R_M which is a modular category with S -matrix S and twists T_i , $i \in I$.

Date: January 23, 2009.

¹We note that our definition of a modular datum is slightly less restrictive than the definition of a modular datum in [G] (where positivity of dimensions is assumed), and the definition of a fusion datum in [L] (where there is an additional involution on the set of labels).

Theorem 1.4. (i) Let $D = \sum_i d_i^2$. Then $D \neq 0$, and one has $S^2 = DC$, where $C_{ij} = \delta_{ij^*}$

(ii) The numbers S_{ij}/d_j (in particular, d_i) and D/d_i^2 are algebraic integers.

(iii) d_i are totally real and $D - 1$ is totally positive. The matrix S/\sqrt{D} (well defined up to sign) is totally unitary.

Proof. (i) By (2), S is invertible. So we have $\delta_j = s_{0j} \sum_i (S^{-1})_{ji} X_i$. Thus for $j \neq m$ we have

$$\sum_{i,r} (S^{-1})_{ji} (S^{-1})_{mr} X_i X_r = 0.$$

In particular, applying η to this equality and using the symmetry of S , we get

$$\sum_i (S^{-1})_{ji} (S_{i^*m}^{-1}) = 0.$$

Thus, $S^{-1}CS^{-1}$ is a diagonal matrix, hence so is SCS .

Now, let N_{ij}^m be the structure constants of R_M . Then by definition

$$S_{ir}S_{jr} = \sum N_{ij}^m S_{mr}S_{0r}.$$

So since $S_{rj} = S_{jr}$ and $S_{0r} = S_{r0} = S_{r^*0}$, we get

$$S_{ir}S_{rj} = \sum N_{ij}^m S_{mr}S_{r^*0}.$$

Summing this over r , and using that SCS is diagonal, we get

$$(S^2)_{ij} = \delta_{ij^*} D.$$

(ii) By definition, S_{ij}/d_j is an eigenvalue of the integer matrix N_i of multiplication by X_i in R_M , thus it is an algebraic integer. Now,

$$\frac{D}{d_i^2} = \sum_j \frac{s_{ij}}{d_i} \frac{s_{ji^*}}{d_i^*},$$

so it is also an algebraic integer.

(iii) $2d_i = d_i + d_{i^*}$ is an eigenvalue of the matrix $N_i + N_{i^*}$, which is a symmetric integer matrix. So d_i is totally real. Next, $(S_{ij} + S_{i^*j})/d_j$ is also an eigenvalue of $N_i + N_{i^*}$, so it is also totally real and hence S_{i^*j} is totally complex conjugate to S_{ij} . This together with (i) implies that under any complex embedding of $\mathbb{Q}(S_{ij})$, $SS^\dagger = D$, where \dagger is the Hermitian adjoint to S , as desired. \square

2. THE ANDERSON-MOORE-VAFA PROPERTY

Let \mathcal{C} be a modular tensor category with simple objects X_i . Consider the space $E_{ijm}^r = \text{Hom}(X_r, X_i \otimes X_j \otimes X_m)$. Using the braiding this space is identified with E_{jim}^r . The hexagon relation for the braiding implies that

$$\det(\beta_{ij,m}^2|_{E_{ijm}^r}) = \det(\beta_{jm}^2|_{E_{ijm}^r}) \det(\beta_{im}^2|_{E_{jim}^r}).$$

Recall that β^2 acts on $\text{Hom}(X_s, X_p \otimes X_q)$ by the scalar $T_p T_q / T_s$. Therefore, the last identity can be rewritten in the form

$$\prod_p (T_p T_m / T_r)^{N_{ij}^p N_{pm}^r} = \prod_p (T_i T_m / T_p)^{N_{im}^p N_{jp}^r} \prod_p (T_j T_m / T_p)^{N_{jm}^p N_{ip}^r}.$$

This implies the following result.

Theorem 2.1. (*Anderson-Moore-Vafa*) *For a categorifiable modular datum, one has*

$$(T_i T_j T_m T_r)^{N_{ijm}^r} = \prod_p T_p^{N_{ij}^p N_{pm}^r + N_{im}^p N_{jp}^r + N_{jm}^p N_{ip}^r},$$

where $N_{ijm}^r := \sum_p N_{ij}^p N_{pm}^r$.

Corollary 2.2. ([E]) *For a categorifiable modular datum, $T^N = 1$ for some N dividing $D^{5/2}$. Therefore, τ/\sqrt{D} is a root of unity of order dividing $4D^{5/2}|I|$.*

Proof. The Anderson-Moore-Vafa equation can be written as the following equation in $\mathbb{Z} \otimes_{\mathbb{Z}} k^\times$:

$$N_{ijm}^r \otimes T_i T_j T_m T_r = \sum_p (N_{ij}^p N_{pm}^r + N_{im}^p N_{jp}^r + N_{jm}^p N_{ip}^r) \otimes T_p.$$

Let R be the ring of integers of the number field generated by d_i . Multiplying the last equation by d_r , and taking the sum over r (remembering that addition in the second component is really multiplication), we get the following equation in $R \otimes_{\mathbb{Z}} k^\times$:

$$d_i d_j d_m \otimes T_i T_j T_m + \sum_r d_r N_{ijm}^r \otimes T_r = \sum_p (N_{ij}^p d_m d_p + N_{im}^p d_j d_p + N_{jm}^p d_i d_p) \otimes T_p.$$

Let us now multiply this equation by d_i and sum over i . Then we get

$$\sum_i d_i^2 d_j d_m \otimes T_i + D d_j d_m \otimes T_j T_m + \sum_r d_j d_m d_r^2 \otimes T_r = \sum_p (2d_j d_m d_p^2 + D N_{jm}^p d_p) \otimes T_p.$$

After cancelations we get

$$D d_j d_m \otimes T_j T_m = D \sum_p N_{jm}^p d_p \otimes T_p.$$

Now multiply this by d_j and sum over j . We get

$$D^2 d_m \otimes T_m + \sum_j D d_m d_j^2 \otimes T_j = D d_m \sum_p d_p^2 \otimes T_p$$

After cancellation we get

$$D^2 d_m \otimes T_m = 0,$$

and using that d_m^2 divides D , we find that T_m is a root of unity of order dividing $D^{5/2}$. The second statement is deduced from the first one by taking the determinant of the equation $(ST)^3 = \tau S^2$ and using that $\det(S)^4 = D^{2|I|}$. \square

2.1. The Ising category. As an example consider the Ising category, with simple objects $X_0 = \mathbf{1}$, $X_1 = \chi$ and $X_2 = X$, where $\chi \otimes \chi = \mathbf{1}$, $\chi \otimes X = X \otimes \chi = X$, and $X \otimes X = \mathbf{1} \oplus \chi$. In this case, we have $s_{01} = s_{11} = 1$, $s_{02} = \sqrt{2}$, $s_{12} = -\sqrt{2}$, and $s_{22} = 0$. In this case, it is easy to check that we must put $T_1 = -1$, but the axioms of a modular datum do not impose any conditions on $\theta = T_2$. On the other hand, the Anderson-Moore-Vafa identity for $i = j = m = r = 2$ gives $\theta^8 = -1$, i.e. θ must be a primitive 16-th root of unity. Thus the Anderson-Moore-Vafa identity does not follow from the axioms of a modular datum. On the other hand, if $\theta^8 = -1$, the corresponding modular datum is categorifiable. (There are eight nondegenerate braided categories with such a fusion ring, and sixteen modular ones, parametrized by choices of $\sqrt{2}$ and by θ ; namely, each braided category has two modular structures, differing by sign of T_2).

REFERENCES

- [E] On Vafa's theorem for tensor categories, P. Etingof, Mathematical research letters, Vol. 9(5), 2002 , pages. 651-658
- [G] T. Gannon Modular data: the algebraic combinatorics of conformal field theory, math/0103044.
- [L] G. Lusztig, Exotic Fourier transform, Duke Math. J. Volume 73, Number 1 (1994), 227-241.

P.E.: DEPARTMENT OF MATHEMATICS, MASSACHUSETTS INSTITUTE OF TECHNOLOGY, CAMBRIDGE, MA 02139, USA

E-mail address: `etingof@math.mit.edu`