

# A BRIEF SUMMARY OF THE STATEMENTS OF CLASS FIELD THEORY

BJORN POONEN

## 0. PROFINITE COMPLETIONS OF TOPOLOGICAL GROUPS

Let  $G$  be a topological group. The profinite completion of  $G$  is

$$\widehat{G} := \varprojlim_U \frac{G}{U},$$

where  $U$  ranges over the finite-index open normal subgroups of  $G$ . There is a natural continuous homomorphism  $G \rightarrow \widehat{G}$  through which every other continuous homomorphism from  $G$  to a profinite group factors uniquely. If  $G$  is profinite already, then  $G \rightarrow \widehat{G}$  is an isomorphism.

In general,  $G \rightarrow \widehat{G}$  need not be injective or surjective. Nevertheless, we think of  $G$  as being almost isomorphic to  $\widehat{G}$ : The finite-index open subgroups of  $G$  are in bijection with those of  $\widehat{G}$ . And finite-index open subgroups of certain Galois groups are what we are interested in...

## 1. LOCAL CLASS FIELD THEORY

### 1.1. Notation associated to a discrete valuation ring.

$\mathcal{O}$ : a complete discrete valuation ring

$K := \text{Frac}(\mathcal{O})$

$v$ : the valuation  $K^\times \rightarrow \mathbb{Z}$

$\mathfrak{p}$ : the maximal ideal of  $\mathcal{O}$

$k$ : the residue field  $\mathcal{O}/\mathfrak{p}$

$K^s$ : a fixed separable closure of  $K$

$K^{\text{ab}}$ : the maximal abelian extension of  $K$  in  $K^s$

$K^{\text{unr}}$ : the maximal unramified extension of  $K$  in  $K^s$

$k^s$ : the residue field of  $K^{\text{unr}}$ , so  $k^s$  is a separable closure of  $k$ .

Equip  $K$  and its subsets with the topology coming from the absolute value  $|x| := \exp(-v(x))$ .

---

*Date:* February 17, 2012.

I thank David Zywinia for comments. The writing of this article was supported by NSF grant DMS-0301280.

## 1.2. Local fields.

**Definition 1.1.** A nonarchimedean local field is a complete discrete-valued field  $K$  as in Section 1.1 such that the residue field  $k$  is finite. An archimedean local field is  $\mathbb{R}$  or  $\mathbb{C}$ .

Facts:

- A nonarchimedean local field of characteristic 0 is isomorphic to a finite extension of  $\mathbb{Q}_p$ .
- A (nonarchimedean) local field of characteristic  $p > 0$  is isomorphic to  $\mathbb{F}_q((t))$  for some power  $q$  of  $p$ .

**1.3. The local Artin homomorphism.** Let  $K$  be a local field. Local class field theory says that there is a homomorphism

$$\theta: K^\times \rightarrow \text{Gal}(K^{\text{ab}}/K)$$

that is almost an isomorphism. The homomorphism  $\theta$  is called the **local Artin homomorphism**. It cannot be literally an isomorphism, because  $\text{Gal}(K^{\text{ab}}/K)$  is a profinite group, hence compact, while  $K^\times$  is not. What is true is that  $\theta$  induces an isomorphism of topological groups  $\widehat{K^\times} \rightarrow \text{Gal}(K^{\text{ab}}/K)$ .

If  $K$  is archimedean, then  $\theta: K^\times \rightarrow \text{Gal}(K^{\text{ab}}/K)$  is surjective and its kernel is the connected component of the identity in  $K^\times$ .

For the rest of Section 1.3, we assume that  $K$  is nonarchimedean. Then  $\theta$  is injective: The choice of a uniformizer  $\pi \in \mathcal{O}$  lets us write  $K^\times = \mathcal{O}^\times \pi^\mathbb{Z} \simeq \mathcal{O}^\times \times \mathbb{Z}$ , and  $\mathcal{O}^\times$  is already profinite, so  $\widehat{K^\times} \simeq \mathcal{O}^\times \times \widehat{\mathbb{Z}}$ . Thus local class field theory says that there is an isomorphism

$$\mathcal{O}^\times \times \widehat{\mathbb{Z}} \rightarrow \text{Gal}(K^{\text{ab}}/K).$$

More canonically, without choosing  $\pi$ , the two horizontal exact sequences below are almost isomorphic:

$$(1) \quad \begin{array}{ccccccc} 0 & \longrightarrow & \mathcal{O}^\times & \longrightarrow & K^\times & \xrightarrow{v} & \mathbb{Z} \longrightarrow 0 \\ & & \parallel & & \downarrow \theta & & \downarrow \\ 0 & \longrightarrow & \text{Gal}(K^{\text{ab}}/K^{\text{unr}}) & \longrightarrow & \text{Gal}(K^{\text{ab}}/K) & \xrightarrow{\text{res}} & \text{Gal}(K^{\text{unr}}/K) \longrightarrow 0 \end{array}$$

With the identification of the group at lower right

$$\text{Gal}(K^{\text{unr}}/K) \simeq \text{Gal}(k^s/k) \simeq \widehat{\mathbb{Z}}$$

mapping the Frobenius automorphism to  $1 \in \widehat{\mathbb{Z}}$ , the right vertical map in (1) becomes the natural inclusion  $\mathbb{Z} \hookrightarrow \widehat{\mathbb{Z}}$ . In other words,  $\theta$  maps  $K^\times$  isomorphically to the set of  $\sigma \in \text{Gal}(K^{\text{ab}}/K)$  inducing an *integer* power of Frobenius on the residue field (as opposed to a  $\widehat{\mathbb{Z}}$ -power). The bottom row of (1) is simply the profinite completion of the top row.

Also from (1), one sees that  $\theta(\mathcal{O}^\times)$  is the inertia subgroup  $\text{Gal}(K^{\text{ab}}/K^{\text{unr}})$  of  $\text{Gal}(K^{\text{ab}}/K)$ , and that  $\theta$  maps any uniformizer to a Frobenius automorphism in  $\text{Gal}(K^{\text{ab}}/K)$ . Moreover, the descending chain

$$\mathcal{O}^\times \supset 1 + \mathfrak{p} \supset 1 + \mathfrak{p}^2 \supset \dots$$

is mapped isomorphically by  $\theta$  to the descending chain of ramification subgroups of  $\text{Gal}(K^{\text{ab}}/K)$  in the upper numbering.

1.4. **Functoriality.** Let  $L$  be a finite extension of  $K$ . Let  $N_{L/K}: L^\times \rightarrow K^\times$  be the norm map. Let  $\theta_L, \theta_K$  be the local Artin homomorphisms associated to  $L, K$ , respectively. Let  $\text{res}: \text{Gal}(L^{\text{ab}}/L) \rightarrow \text{Gal}(K^{\text{ab}}/K)$  be the homomorphism mapping an automorphism  $\sigma$  of  $L^{\text{ab}}$  to its restriction  $\sigma|_{K^{\text{ab}}}$ . Then the square

$$\begin{array}{ccc} L^\times & \xrightarrow{\theta_L} & \text{Gal}(L^{\text{ab}}/L) \\ N_{L/K} \downarrow & & \downarrow \text{res} \\ K^\times & \xrightarrow{\theta_K} & \text{Gal}(K^{\text{ab}}/K) \end{array}$$

commutes.

1.5. **Finite abelian extensions.** Because  $\theta$  is almost an isomorphism, and because of Galois theory, the following sets are in bijection:

- The finite-index open subgroups of  $K^\times$ .
- The (finite-index) open subgroups of  $\text{Gal}(K^{\text{ab}}/K)$ .
- The finite abelian extensions of  $K$  contained in  $K^{\text{s}}$ .

Going backwards, if  $L$  is a finite abelian extension of  $K$  in  $K^{\text{s}}$ , the corresponding subgroup of  $K^\times$  is  $N_{L/K}L^\times$ . (This could be viewed as a consequence of the functoriality above.)

The composition

$$K^\times \rightarrow \text{Gal}(K^{\text{ab}}/K) \xrightarrow{\text{res}} \text{Gal}(L/K)$$

is surjective with kernel  $N_{L/K}L^\times$ , and  $\mathcal{O}^\times$  maps to the inertia subgroup  $I_{L/K} \trianglelefteq \text{Gal}(L/K)$ , and any uniformizer  $\pi$  maps to a Frobenius element of  $\text{Gal}(L/K)$ .

## 2. GLOBAL CLASS FIELD THEORY (VIA IDELES)

### 2.1. Global fields.

**Definition 2.1.** A number field is a finite extension of  $\mathbb{Q}$ . A global function field is a finite extension of  $\mathbb{F}_p(t)$  for some prime  $p$ , or equivalently is the function field of a geometrically integral curve over a finite field  $\mathbb{F}_q$  (called the **constant field**), where  $q$  is a power of some prime  $p$ . A global field is a number field or a global function field.

Throughout Sections 2 and 3,  $K$  is a global field. If  $v$  is a nontrivial place of  $K$  (given by an absolute value on  $K$ ), then the completion  $K_v$  is a local field. If  $v$  is nonarchimedean, let  $\mathcal{O}_v$  be the valuation subring of  $K_v$ ; if  $v$  is archimedean, let  $\mathcal{O}_v = K_v$ .

2.2. **The adèle ring.** The adèle ring of  $K$  is the restricted direct product

$$\mathbf{A}_K := \prod'_v (K_v, \mathcal{O}_v) := \left\{ (a_v) \in \prod_v K_v : a_v \in \mathcal{O}_v \text{ for all but finitely many } v \right\}.$$

It is a topological ring: the topology is uniquely characterized by the condition that  $\prod_v \mathcal{O}_v$  is open in  $\mathbf{A}_K$  and has the product topology. The diagonal map  $K \rightarrow \mathbf{A}_K$  is like  $\mathbb{Z} \rightarrow \mathbb{R}$ : it embeds  $K$  as a discrete co-compact subgroup of  $\mathbf{A}_K$ .

2.3. **The idele group and idele class group.** The idele group of  $K$  is

$$\mathbf{A}_K^\times = \prod'_v (K_v^\times, \mathcal{O}_v^\times) := \left\{ (a_v) \in \prod_v K_v^\times : a_v \in \mathcal{O}_v^\times \text{ for all but finitely many } v \right\}.$$

It is a topological group: the topology is uniquely characterized by the condition that  $\prod_v \mathcal{O}_v^\times$  is open in  $\mathbf{A}_K^\times$  and has the product topology.<sup>1</sup> The diagonal map  $K^\times \rightarrow \mathbf{A}_K^\times$  is like  $\mathbb{Z}^\times \rightarrow \mathbb{R}^\times$ : it embeds  $K^\times$  as a discrete subgroup of  $\mathbf{A}_K^\times$ , but the quotient  $C_K := \mathbf{A}_K^\times / K^\times$  is not compact. The topological group  $C_K$  is called the **idele class group**.

2.4. **The global Artin homomorphism.** Let  $K^s$  be a fixed separable closure of  $K$ . Let  $K^{\text{ab}}$  be the maximal abelian extension of  $K$  contained in  $K^s$ .

The group  $C_K$  plays the role in global class field theory played by  $K^\times$  in local class field theory. Namely, if  $K$  is a global field, there is a **global Artin homomorphism**

$$\theta: C_K \rightarrow \text{Gal}(K^{\text{ab}}/K)$$

that induces an isomorphism  $\widehat{C}_K \xrightarrow{\sim} \text{Gal}(K^{\text{ab}}/K)$ .

If  $K$  is a number field, then  $\theta$  is surjective and its kernel is the connected component of the identity in  $C_K$ .

If  $K$  is a global function field with constant field  $k$ , then  $\theta$  is injective and  $\theta(C_K)$  equals the set of  $\sigma \in \text{Gal}(K^{\text{ab}}/K)$  whose restriction in  $\text{Gal}(k^s/k)$  is an *integer* power of the Frobenius generator.

2.5. **Functoriality.** Let  $L$  be a finite extension of  $K$  of degree  $n$ . Then  $\mathbf{A}_L \simeq \mathbf{A}_K \otimes_K L$  is free of rank  $n$  over  $\mathbf{A}_K$ , so there is a norm map  $N_{L/K}: \mathbf{A}_L \rightarrow \mathbf{A}_K$ . We write  $N_{L/K}$  also for the induced homomorphism  $N_{L/K}: C_L \rightarrow C_K$ . Then

$$\begin{array}{ccc} C_L & \xrightarrow{\theta_L} & \text{Gal}(L^{\text{ab}}/L) \\ N_{L/K} \downarrow & & \downarrow \text{res} \\ C_K & \xrightarrow{\theta_K} & \text{Gal}(K^{\text{ab}}/K) \end{array}$$

commutes.

2.6. **Finite abelian extensions.** The following sets are in bijection:

- The finite-index open subgroups of  $C_K$ .
- The finite-index open subgroups of  $\text{Gal}(K^{\text{ab}}/K)$ .
- The finite abelian extensions of  $K$  contained in  $K^s$ .

Going backwards, if  $L$  is a finite abelian extension of  $K$  in  $K^s$ , the corresponding subgroup of  $C_K$  is  $N_{L/K}C_L$ . The composition

$$C_K \rightarrow \text{Gal}(K^{\text{ab}}/K) \xrightarrow{\text{res}} \text{Gal}(L/K)$$

is surjective with kernel  $N_{L/K}C_L$ .

<sup>1</sup>Alternatively, one can use the general recipe for getting the topology on the units of a topological ring  $R$ : not the subspace topology on  $R^\times$  as a subset of  $R$  (this may fail to make the inverse map  $R^\times \rightarrow R^\times$  continuous), but the subspace topology on the set of solutions to  $xy = 1$  in  $R \times R$  (this is what one gets if one expresses the multiplicative group scheme  $\mathbb{G}_m$  as an affine variety).

**2.7. Connection between the global and local Artin homomorphisms.** Let  $v$  be a place of  $K$ . Identify  $K_v^\times$  with a subgroup of  $\mathbf{A}_K^\times$  by mapping  $\alpha \in K_v^\times$  to the idele with  $\alpha$  in the  $v$ -th position and 1 in every other position. The composition  $K_v^\times \hookrightarrow \mathbf{A}_K^\times \twoheadrightarrow C_K$  is injective. Let  $\theta_v$  be the local Artin homomorphism for  $K_v$ . Then the diagram

$$\begin{array}{ccc} K_v^\times & \xrightarrow{\theta_v} & \text{Gal}(K_v^{\text{ab}}/K_v) \\ \downarrow & & \downarrow \text{res} \\ C_K & \xrightarrow{\theta} & \text{Gal}(K^{\text{ab}}/K) \end{array}$$

commutes. Thus  $\theta$  determines  $\theta_v$ .

Conversely, if one knows  $\theta_v$  for all  $v$ , one can construct  $\theta$  as follows. Let  $L$  be a finite abelian extension of  $K$  contained in  $K^{\text{s}}$ . Define

$$\begin{aligned} \mathbf{A}_K^\times &\rightarrow \text{Gal}(L/K) \\ (a_v) &\mapsto \prod_v \theta_v(a_v); \end{aligned}$$

if  $v$  is unramified in  $L/K$ , and  $a_v \in \mathcal{O}_v^\times$ , then  $\theta_v(a_v) = 1$ , so all but finitely many terms in the infinite product are 1, and the product makes sense. Take the inverse limit over all possible  $L$  to get

$$\mathbf{A}_K^\times \rightarrow \text{Gal}(K^{\text{ab}}/K).$$

The idelic version of the Artin reciprocity law says that  $K^\times$  is in the kernel, so we get a homomorphism

$$C_K \rightarrow \text{Gal}(K^{\text{ab}}/K),$$

which is  $\theta$ .

## 2.8. Moduli.

**Definition 2.2.** A **modulus** is a formal product  $\mathfrak{m} = \prod_v v^{e_v}$  where  $e_v \in \mathbb{Z}_{\geq 0}$ , all but finitely many  $e_v$  equal 0, and  $e_v \in \{0, 1\}$  for real  $v$ , and  $e_v = 0$  for complex  $v$ . The **support**  $\text{supp } \mathfrak{m}$  is the (finite) set of *nonarchimedean* places  $v$  such that  $e_v \neq 0$ .

If  $K$  is a number field, then a modulus can be viewed as a pair consisting of

- (1) an integral ideal of the ring of integers  $\mathcal{O}_K$ , and
- (2) a subset of the real places.

If  $K$  is the function field of a smooth projective curve  $X$  over a finite field, then a modulus is the same thing as an effective divisor on  $X$ .

**2.9. Ray class groups and ray class fields.** In this section we assume that  $K$  is a number field. Fix a modulus  $\mathfrak{m} = \prod_v v^{e_v}$ . We will define a finite-index open subgroup  $U_{\mathfrak{m},v} \subseteq \mathcal{O}_v^\times$  for each  $v$ . If  $e_v = 0$ , define  $U_{\mathfrak{m},v} := \mathcal{O}_v^\times$ . If  $e_v > 0$  and  $v$  is nonarchimedean, define  $U_{\mathfrak{m},v} := 1 + \mathfrak{p}_v^{e_v}$ , where  $\mathfrak{p}_v$  is the maximal ideal of  $\mathcal{O}_v$ . If  $e_v > 0$  and  $v$  is real, define  $U_{\mathfrak{m},v}$  as  $\mathbb{R}_{>0} \subseteq \mathbb{R}^\times \simeq K_v^\times$ . Define  $U_{\mathfrak{m}} := \prod_v U_{\mathfrak{m},v} \subseteq \mathbf{A}_K^\times$ . The image of  $U_{\mathfrak{m}}$  under  $\mathbf{A}_K^\times \twoheadrightarrow C_K$  is a finite-index open subgroup  $U'_{\mathfrak{m}}$  of  $C_K$  (this is equivalent to finiteness of the class number of  $K$ , as we will see in Section 3.4). The corresponding finite abelian extension  $R_{\mathfrak{m}}$  of  $K$  is

called the ray class field of modulus  $\mathfrak{m}$ , and  $R_{\mathfrak{m}}$  over  $K$  is unramified at all  $v$  with  $e_v = 0$ . The ray class group of modulus  $\mathfrak{m}$  is

$$\frac{C_K}{U'_{\mathfrak{m}}} = \frac{\mathbf{A}_K^{\times}}{U_{\mathfrak{m}}K^{\times}},$$

which is isomorphic to  $\text{Gal}(R_{\mathfrak{m}}/K)$  via the global Artin homomorphism.

Every finite-index open subgroup of  $\mathbf{A}_K^{\times}$  contains  $U_{\mathfrak{m}}$  for some  $\mathfrak{m}$ , so every finite abelian extension of  $K$  is contained in  $R_{\mathfrak{m}}$  for some  $\mathfrak{m}$ .

### 3. GLOBAL CLASS FIELD THEORY (VIA IDEALS)

In this section we assume that  $K$  is a number field.

**3.1. Classical ray class groups.** Let  $I$  be the group of fractional ideals of  $K$ , or equivalently, the free abelian group on the nonarchimedean places of  $K$ . Let  $P$  be the subgroup of principal ideals. The class group is  $\text{Cl } \mathcal{O}_K := I/P$ .

We now generalize to an arbitrary modulus  $\mathfrak{m} = \prod_v v^{e_v}$ . Let  $I_{\mathfrak{m}}$  be the subgroup of fractional ideals that do not involve the primes dividing  $\mathfrak{m}$ ; i.e.,  $I_{\mathfrak{m}}$  is the free abelian group on the nonarchimedean places  $v$  satisfying  $e_v = 0$ . For  $a \in K^{\times}$ , the notation  $a \equiv 1 \pmod{\times \mathfrak{m}}$  means that  $a \in U_{\mathfrak{m},v}$  for every  $v$  satisfying  $e_v > 0$ . The group  $P_{\mathfrak{m}} \subseteq I_{\mathfrak{m}}$  is the group of principal ideals generated by some  $a \in K^{\times}$  with  $a \equiv 1 \pmod{\times \mathfrak{m}}$ . The classical ray class group of modulus  $\mathfrak{m}$  is  $\text{Cl}_{\mathfrak{m}} \mathcal{O}_K := I_{\mathfrak{m}}/P_{\mathfrak{m}}$ . Section 3.4 will prove that this is isomorphic to the ray class group  $C_K/U'_{\mathfrak{m}}$  defined in Section 2.9.

**3.2. The classical Artin homomorphism.** Let  $L/K$  be a finite abelian extension of number fields. Let  $S$  be a finite set of finite primes of  $K$  such that  $S$  contains every prime that ramifies in  $L$ . Let  $I_S$  be the group of fractional ideals that do not involve the primes in  $S$ . The classical Artin homomorphism is the map

$$\Theta: I_S \rightarrow \text{Gal}(L/K)$$

sending each prime ideal  $\mathfrak{p} \notin S$  to the Frobenius element  $\text{Frob}_{\mathfrak{p}} \in \text{Gal}(L/K)$ .

**3.3. The main theorems.** The Artin reciprocity law states that there exists a modulus  $\mathfrak{m}$  (depending on  $L/K$ ) with  $\text{supp } \mathfrak{m} = S$  such that the subgroup  $P_{\mathfrak{m}} \subseteq I_{\mathfrak{m}} = I_S$  is contained in  $\ker \Theta$ . The existence theorem states that given a modulus  $\mathfrak{m}$  and group  $H$  with  $P_{\mathfrak{m}} \subseteq H \subseteq I_{\mathfrak{m}}$  there exists an abelian extension  $L$  of  $K$  unramified outside  $\text{supp } \mathfrak{m}$  such that the kernel of  $\Theta$  for  $L/K$  equals  $H$ .

**3.4. Comparison of ideal groups and idele groups.** Consider the trivial modulus  $\mathfrak{m} = 1$  (with  $e_v = 0$  for all  $v$ ). Taking the restricted direct product of the valuation maps  $v: K_v^{\times} \rightarrow \mathbb{Z}$  gives a surjective homomorphism

$$\mathbf{A}_K^{\times} \rightarrow I$$

that discards the archimedean components of its input, and its kernel is  $U_1 = \prod_v \mathcal{O}_v^{\times}$ . Thus  $\frac{\mathbf{A}_K^{\times}}{U_1} \simeq I$ . If we take the quotient by the image of  $K^{\times}$  on both sides, we find that the ray class group  $\frac{\mathbf{A}_K^{\times}}{U_1 K^{\times}}$  of modulus 1 is isomorphic to the class group  $I/P = \text{Cl } \mathcal{O}_K$ . The ray class field  $R_1$  of modulus 1 is called the Hilbert class field, which can be characterized also as the

maximal abelian extension of  $K$  in  $K^s$  that is unramified at all places of  $K$  (including the archimedean ones). We get

$$\frac{C_K}{U'_1} = \frac{\mathbf{A}_K^\times}{U_1 K^\times} \simeq \frac{I}{P} = \text{Cl } \mathcal{O}_K \simeq \text{Gal}(R_1/K).$$

This can be generalized to an arbitrary modulus  $\mathfrak{m} = \prod v^{e_v}$  as follows. Let  $\mathbf{A}_K^{\mathfrak{m}} \subseteq \mathbf{A}_K^\times$  be the subgroup consisting of  $(a_v)$  with  $a_v \in U_{\mathfrak{m},v}$  for every  $v$  with  $e_v > 0$ . Let  $K^{\mathfrak{m}} = \mathbf{A}_K^{\mathfrak{m}} \cap K^\times$ . We have an isomorphism

$$\frac{\mathbf{A}_K^{\mathfrak{m}}}{U_{\mathfrak{m}}} \xrightarrow{\sim} I_{\mathfrak{m}}.$$

Dividing by the image of  $K^{\mathfrak{m}}$  on both sides gives

$$(2) \quad \frac{\mathbf{A}_K^{\mathfrak{m}}}{U_{\mathfrak{m}} K^{\mathfrak{m}}} \xrightarrow{\sim} \frac{I_{\mathfrak{m}}}{P_{\mathfrak{m}}}.$$

On the other hand,  $\mathbf{A}_K^\times = \mathbf{A}_K^{\mathfrak{m}} K^\times$ , so there is an isomorphism

$$\frac{\mathbf{A}_K^{\mathfrak{m}}}{K^{\mathfrak{m}}} \xrightarrow{\sim} \frac{\mathbf{A}_K^\times}{K^\times} = C_K.$$

Dividing by the image of  $U_{\mathfrak{m}}$  on both sides, and combining with (2), we get isomorphisms

$$\frac{C_K}{U'_m} = \frac{\mathbf{A}_K^\times}{U_m K^\times} \simeq \frac{I_m}{P_m} = \text{Cl}_m \mathcal{O}_K \simeq \text{Gal}(R_m/K).$$

#### 4. AN INTRODUCTION TO AN INTRODUCTION TO THE LANGLANDS PROGRAM

Let  $K$  be a local or global field. Every 1-dimensional character (continuous homomorphism)

$$\text{Gal}(K^s/K) \rightarrow \mathbb{C}^\times$$

factors through  $\text{Gal}(K^{\text{ab}}/K)$  and has finite image. These characters form a discrete abelian group, the Pontryagin dual of the profinite group  $\text{Gal}(K^{\text{ab}}/K)$ . It follows that the problem of classifying finite abelian extensions of  $K$  is more or less the same as the problem of describing all these characters.

The Langlands program is an attempt to understand  $\text{Gal}(K^s/K)$  more completely by describing its higher-dimensional representations: the group  $\mathbb{C}^\times = \text{GL}_1(\mathbb{C})$  is replaced by  $\text{GL}_n(\mathbb{C})$ , or even  $G(\mathbb{C})$  for other linear algebraic groups  $G$ . The continuous homomorphisms

$$\text{Gal}(K^s/K) \rightarrow G(\mathbb{C})$$

are conjectured to correspond to certain ‘‘automorphic’’ objects defined intrinsically in terms of  $K$ , just as class field theory gives a description of the group  $\text{Gal}(K^{\text{ab}}/K)$  (which is defined in terms of extrinsic objects such as finite abelian extensions, which are initially mysterious) in terms of intrinsic objects ( $K^\times$  or  $C_K$ ) obtained directly from  $K$ .

Ultimately, the program would give information about nonabelian algebraic extensions of  $K$ .

## 5. FURTHER READING

For basics on profinite groups, see [Ser02, I.§1] and [Gru86]. The latter discusses infinite Galois theory as well.

For local class field theory, see [Ser86]. For the approach to global class field theory via cohomology of ideles, see [Tat86]. For a treatment of global class field theory via ideals, see [Jan96]. All these topics are covered also in [Neu99].

For an introduction to the Langlands program, see [BG03].

## REFERENCES

- [BG03] D. Bump, J. W. Cogdell, E. de Shalit, D. Gaitsgory, E. Kowalski, and S. S. Kudla, *An introduction to the Langlands program*, Birkhäuser Boston Inc., Boston, MA, 2003. Lectures presented at the Hebrew University of Jerusalem, Jerusalem, March 12–16, 2001; Edited by Joseph Bernstein and Stephen Gelbart. MR1990371 (2004g:11037) ↑5
- [Gru86] K. Gruenberg, *Profinite groups*, Algebraic number theory (J. W. S. Cassels and A. Fröhlich, eds.), Academic Press Inc. [Harcourt Brace Jovanovich Publishers], London. Reprint of the 1967 original, 1986, pp. 116–127. MR0225922 (37 #1512) ↑5
- [Jan96] Gerald J. Janusz, *Algebraic number fields*, 2nd ed., Graduate Studies in Mathematics, vol. 7, American Mathematical Society, Providence, RI, 1996. MR1362545 (96j:11137) ↑5
- [Neu99] Jürgen Neukirch, *Algebraic number theory*, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 322, Springer-Verlag, Berlin, 1999. Translated from the 1992 German original and with a note by Norbert Schappacher; With a foreword by G. Harder. MR1697859 (2000m:11104) ↑5
- [Ser86] J.-P. Serre, *Local class field theory*, Algebraic number theory (J. W. S. Cassels and A. Fröhlich, eds.), Academic Press Inc. [Harcourt Brace Jovanovich Publishers], London. Reprint of the 1967 original, 1986, pp. 128–161. MR0220701 (36 #3753) ↑5
- [Ser02] Jean-Pierre Serre, *Galois cohomology*, Corrected reprint of the 1997 English edition, Springer Monographs in Mathematics, Springer-Verlag, Berlin, 2002. Translated from the French by Patrick Ion and revised by the author. MR1867431 (2002i:12004) ↑5
- [Tat86] J. T. Tate, *Global class field theory*, Algebraic number theory (J. W. S. Cassels and A. Fröhlich, eds.), Academic Press Inc. [Harcourt Brace Jovanovich Publishers], London. Reprint of the 1967 original, 1986, pp. 162–203. MR0220697 (36 #3749) ↑5

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, BERKELEY, CA 94720-3840, USA  
*E-mail address:* `poonen@math.berkeley.edu`  
*URL:* `http://math.berkeley.edu/~poonen`