

LECTURE 2

Iterated blow ups and manifolds with corners

Last time I went through the definition of the manifold $[M; Y]$ obtained from M by blowing up along a closed embedded submanifold Y with its natural blow-down map

$$(2.1) \quad \beta : [M; Y] \longrightarrow M.$$

This is a smooth map, so pull-back gives $\beta^* : \mathcal{C}^\infty(M) \longrightarrow \mathcal{C}^\infty([M; Y])$. This is injective but cannot be surjective, namely there are more functions which are smooth in polar coordinates. This in fact is one of the reasons to blow up.

What about vector fields? The vector fields which lift to be smooth under β are precisely those which are tangent to Y . There are always local coordinates z, y in M near any point of Y in which Y is locally defined by the $z_1 = \dots = z_k = 0$ and the y_i 's become coordinates on Y . The vector fields tangent to Y are then the \mathcal{C}^∞ combinations of the ∂_{y_i} and the $z_j \partial_{z_i}$. The approach I took last time shows that these lift to be smooth, to be tangent to the new boundary $r = 0$ and to span, over \mathcal{C}^∞ coefficients on $[M; Y]$ all the vector fields tangent to $\partial[M; Y]$. For any manifold with boundary this latter space consists of all the sections of a vector bundle

$$(2.2) \quad \{V \in \mathcal{C}^\infty(X; TX); V \text{ is tangent to } \partial X\} = \mathcal{C}^\infty(X; {}^bTX).$$

This is already an important fact, since a Lie algebra of vector fields consisting of all the smooth sections of a vector bundle is getting close to the standard case of all the smooth vector fields on a (compact) manifold without boundary.

1. Manifolds with corners

Each blow up introduces a boundary, so in order to do iterated blow up we have to work in the context of manifolds with corners. I will be brief about these, really there is not much to worry about in the basic theory. In summary the definition of a manifold in the usual sense is as a set X with a covering by local coordinates systems with \mathcal{C}^∞ transition maps. For a manifold with corners we allow the coordinate 'model' to be the intersection of an open subset of \mathbb{R}^n with one of the k -corners $[0, \infty)^k \times \mathbb{R}^{n-k}$. Smoothness of a map is the existence of a smooth extension to an open set in \mathbb{R}^n , by Whitney's (easy) extension theorem this is the same as local boundedness of all derivatives. So, locally a manifold with corners looks like $[0, \infty)^k \times \mathbb{R}^{n-k}$ at a boundary point of codimension k ; I will write the local coordinates $x_1, \dots, x_k, y_1, \dots, y_{n-k}$. This means that there are boundary hypersurfaces, connected sets given locally by the vanishing of one of the x_j . To make sure that these are manifolds with corners in the same sense I insist that the boundary hypersurfaces be embedded. This means that each of them, $H \subset X$ is given by $\rho_H = 0$ where $\rho_H \in \mathcal{C}^\infty(X)$, $d\rho_H \neq 0$ on H and $\rho_H \geq 0$.

So, as a little exercise you can go back to what I did last time and see that we can blow up closed ‘embedded’ submanifolds $Y \subset X$ for a manifold with corners, provided that embedded implies that an appropriate version of the collar neighbourhood theorem holds. This is the condition that H be a p-submanifold. More precisely this means that near each point of X there are local coordinates of the adapted sort that I described above, x and y such that locally

$$(2.3) \quad Y = \{x_1 = \cdots = x_j = 0, y_i = \cdots = y_l = 0\}$$

where either $j = 0$ (no x equations) or $l = 0$ (no y equations) is permitted. Note that $l = 0$ makes Y into a boundary face – a component of the intersection of boundary hypersurfaces. The other extreme $j = 0$ corresponds to an ‘interior p-submanifold’ which is most like the usual case.

PROPOSITION 3. *It is always possible to blow up a closed embedded p-submanifold Y in a manifold with corners X giving a new manifold with corners $[X; Y]$ with maximal boundary codimension either the same or increased by one and with a smooth blow down map*

$$(2.4) \quad \beta : [X; Y] \longrightarrow X.$$

Although there is nothing much to manifolds with corners at the level I have described here, there is something more significant in the maps between them which I want to emphasize. Smoothness itself is straightforward, but smooth maps between manifolds with corners $f : X \longrightarrow X'$, can, and should, be required to ‘preserve some of the boundary structure’. The natural condition is that inverse images of boundary faces should be boundary faces, in a wide sense that they be unions of boundary faces. In terms of boundary defining functions this means

$$(2.5) \quad f^* \rho'_i = a_i \prod_j \rho_j^{\alpha_{ij}}, \quad 0 < a_i \in \mathcal{C}^\infty(X)$$

where the $\rho'_i \in \mathcal{C}^\infty(X')$ and $\rho_j \in \mathcal{C}^\infty(X)$ are listings of the boundary defining functions for X' and X respectively; the α_{ij} are necessarily non-negative integers but 0 is allowed.

DEFINITION 1. A smooth map $f : X \longrightarrow X'$ between manifolds with corners which satisfies (2.5) is called an *interior b-map*.

Here the ‘b-’ just stands for boundary. Note that the composite of two b-maps is again a b-map. A general, not necessarily ‘interior’ b-map is one which is an interior b-map into one of the boundary faces of X' . This just corresponds to either (2.5) or $f^* \rho'_i \equiv 0$ holding for each boundary defining function of X' .

2. Examples again

Now, I can describe one of the original applications of blow up – to define the b-calculus (although this had already been done, maybe it is better to say it gives a clear characterization).

Above, I emphasized p-submanifolds. However, one of the most interesting examples of embedded submanifolds in the usual setting of a compact manifold without boundary is the diagonal

$$(2.6) \quad \text{Diag} = \{(m, m) \in M^2; m \in M\}.$$

This is certainly embedded. However in the case of a manifold with boundary it is not a p-submanifold as we see even in the one-dimensional case

$$(2.7) \quad \text{Diag} = \{(x_1, x_2) \in [0, 1]^2; x_1 = x_2\}.$$

[Sketch]

As we have been hearing from Michael Taylor, pseudodifferential operators correspond to kernels with rather simple ‘conormal’ singularities at the diagonal and smooth elsewhere. In this case there are several possibilities about what to do. We can ignore the boundary, defining pseudodifferential operators by restriction from $\mathbb{R} \times \mathbb{R}$ for instance. However, ignoring boundaries that are really there is not wise. We can follow Boutet de Monvel and consider *transmission conditions* – perhaps Gerd Grubb will talk more about this. However, we can also think of defining pseudodifferential operators instead as generalizations of the *tangent* vector fields

$$(2.8) \quad x\partial_x, \partial_{y_j}.$$

So, this brings us to two questions simultaneously.

- (1) If we think of non-p-submanifolds as singular, how can we resolve the diagonal in the case of a manifold with boundary (or for that matter with corners).
- (2) What does it mean to ‘resolve’ the algebra of vector fields tangent to the boundary on a manifold with boundary (of with corners).

In the first case we can say a blow up of some Y resolves the diagonal if the lift of Diag to $[M^2; Y]$ is a p-submanifold. In the second case we say the blow up resolves the Lie algebra if its elements lift, from one of the factors, to be smooth and also to be collectively transversal to the diagonal.

We already know that for the vector fields to lift to be smooth, they must be tangent to Y . Clearly $Y \supset \partial \text{Diag}$ is also necessary, since otherwise there are points at which nothing is changed and Diag cannot have been resolved. In the case of the tangent vector fields (2.8), these two conditions force

$$(2.9) \quad Y = \partial X \times \partial X.$$

This is a boundary face, and hence a p-submanifold. I am assuming here that ∂X is connected.

LEMMA 5. *The diagonal $\text{Diag} \subset X^2$ for a compact manifold with boundary lifts to $X_b^2 = [X^2; (\partial X)^2]$ to a p-submanifold and $\mathcal{V}_b(X)$, the Lie algebra of smooth vector fields lifts to be transversal to the (lifted) diagonal.*

PROOF. Computation. In fact if you think about it this really reduces to the 1-dimensional case. I have not yet defined the lift of a submanifold under blow up, so you should continue reading to find out what this means. \square

3. Commutation

Now, if Y_1 and Y_2 are both subsets of M we can ask what happens to Y_2 after we blow up Y_1 – which better be a p-submanifold for this to be possible. We have to distinguish between the two cases where $Y_2 \subset Y_1$ and $Y_2 \setminus Y_1 \neq \emptyset$. In the first case we define the lift $\widetilde{Y}_2 = \beta^!(Y_2) \subset [X; Y_1]$ to be $\beta^{-1}(Y_2)$. In the second case we define it to be the closure of $\beta^{-1}(Y_2 \setminus Y_1)$ – although this doesn’t make much sense unless Y_2 meets Y_1 reasonably sensibly.

[Sketch]

It is easy to think of a ‘joint p-submanifold’ condition on $Y_1, Y_2 \subset X$ – namely that they are each p-submanifolds and near any point of their intersections there is *one* adapted coordinate system in M in terms of which the *both* take the form (2.3), with different ‘index sets’ of course so we should generalize this by saying

$$(2.10) \quad Y_p = \{x_i = 0, i \in I_p \subset \{1, \dots, k\}, y_j = 0, j \in J_p \subset \{1, \dots, n - k\}\}.$$

PROPOSITION 4. *If Y_1 and Y_2 are joint p-submanifolds in the sense of (2.10) then the lift of Y_2 to $[X; Y_1]$ is a p-submanifold. In the special case that in addition either $Y_1 \subset Y_2$ or $Y_2 \subset Y_1$ or $Y_1 \pitchfork Y_2$ there is a canonical isomorphism*

$$(2.11) \quad [[X; Y_1]; \beta_1^!(Y_2)] = [[X; Y_2]; \beta_2^!(Y_1)]$$

but not otherwise.

Generally we denote the iterated blow up, $[[X; Y_1]; \beta_1^!(Y_2)]$ as $[X; Y_1; Y_2]$ and then the commutation result becomes

$$(2.12) \quad [X; Y_1; Y_2] = [X; Y_2; Y_1].$$

PROOF. I doubt that I will have time to do this in the lecture but it is not so hard. Note that the transversal case, $Y_1 \pitchfork Y_2$ is the easy one. In terms of (2.10) it means that $I_1 \cap I_2 = \emptyset = J_1 \cap J_2$. What this amounts to is that one can locally decompose $M = M_1 \times M_2$ as a product, so that $Y_1 = Y_1' \times M_2$ and $Y_2 = M_1 \times Y_2'$ where $Y_1' \subset M_1$ and $Y_2' \subset M_2$ are p-submanifolds. Then it follows easily.

The case of inclusion one way or the other can be done by computation. One way to think about it is to consider the radial vector fields around the submanifolds. For Y in (2.3) this would be

$$(2.13) \quad x_1 \partial_{x_1} + \dots + x_j \partial_{x_j} + y_1 \partial_{y_1} + \dots + y_l \partial_{y_l}.$$

The condition of inclusion means that one of these vector fields (the one for the smaller submanifold) is obtained from the other by adding terms. Since this radial vector field lifts to $r\partial_r$, it follows that the radial actions commute and this leads to the commutation of the blow ups.

The fact that this *doesn't* work otherwise can also be seen by lifting the radial vector fields. \square

4. Tangent vector fields again

Let me point out that this commutation result allows us to resolve the Lie algebra of vector fields, $\mathcal{V}_b(X)$, tangent to all the boundaries of a manifold with corners, and hence as I will indicate below, to define the b-calculus in this context too. Namely, for a manifold with corners X consider all the products $H_i \times H_i$ of a boundary hypersurface with itself. These are all transversal one to another. So we need a little result to proceed.

LEMMA 6. *Under blow up of a boundary face all other boundary faces lift to boundary faces and transversal boundary faces remain transversal.*

So, combining this with the commutation result for transversal p-submanifolds discussed above, we can define unambiguously

$$(2.14) \quad X_b^2 = [X^2; H_1^2; H_2^2; \dots; H_N^2]$$

giving a manifold with corners, independent of the order, to which Diag lifts to be a p -submanifold. The the tangent vector fields, forming $\mathcal{V}_b(X)$, lift to be collectively transversal to this lifted diagonal.

5. Another commutation result

In Proposition 4 it is noted that for joint p -submanifolds Y_1 and Y_2 which are neither comparable (meaning one is contained in the other) nor transversal, the two manifolds $[X; Y_1; Y_2]$ and $[X; Y_2; Y_1]$ are different. How then can one ‘blow up’ such a subset. It is possible to show that one can ‘correct’ the blow up in two ways.

The first, and most frequent ‘solution’ is to simply blow up the intersection first and then check that

$$(2.15) \quad [X; Y_1 \cap Y_2; Y_1; Y_2] \equiv [X; Y_1 \cap Y_2; Y_2; Y_1].$$

In fact, after under the blow up of $Y_1 \cap Y_2$ the two bigger manifolds Y_1 and Y_2 lift to p -submanifolds which are disjoint, and hence transversal – giving (2.15).

There is a second alternative, which is rarely used (and may not even be in the literature). That is, one can blow up $Y_1 \cap Y_2$ *last*:

PROPOSITION 5. *For any pair of embedded joint p -submanifolds there is a natural diffeomorphism*

$$(2.16) \quad [X; Y_1; Y_2; Y_1 \cap Y_2] \equiv [X; Y_2; Y_1; Y_1 \cap Y_2].$$

However this manifold is different to the one in (2.15). Note that on the left in (2.16) $Y_1 \cap Y_2$ first lifts as a submanifold of Y_1 but *is not* a submanifold of the lift of Y_2 – so the notation is a bit dangerous.

6. Fibrations and b-fibrations

Perhaps the most important smooth maps between manifolds are diffeomorphisms. However, in geometric and other settings *fibrations* are also of vital importance. A smooth map $f : X \rightarrow X'$ between manifolds without boundary is a *submersion* if its differential is everywhere surjective, $f_* : T_x X \rightarrow T_{f(x)} X'$, for all $x \in X$. If X and X' are compact the Implicit Function Theorem shows that f is actually a fibration, meaning that it is surjective and each point $x' \in X'$ has an open neighbourhood U for which there is a diffeomorphism F giving a commutative diagram

$$(2.17) \quad \begin{array}{ccc} f^{-1}(U) & \xrightarrow{F} & U \times Z \\ f \downarrow & \swarrow \pi_1 & \\ U; & & \end{array}$$

here π_1 is projection onto the first factor. The manifold Z is then determined up to diffeomorphism (provided X' is connected) and such a triple $f : X \rightarrow X'$ may be written

$$(2.18) \quad \begin{array}{ccc} Z & \text{---} & X \\ & & \downarrow f \\ & & X' \end{array}$$

(there is now actual map from the model fibre Z , rather each fibre is diffeomorphic to it) and thought of as a fibre bundle, with fibre Z and structure group $\text{Diff}(Z)$. One reason such maps are particularly well-behaved is that Fubini's theorem shows that fibre-integration preserves smoothness:

$$(2.19) \quad f_* : \mathcal{C}_c^\infty(X; \Omega) \longrightarrow \mathcal{C}_c^\infty(X'; \Omega).$$

Here Ω is the (trivial) bundle of densities, those things which can be invariantly integrated.

We can easily set up fibrations in the category of compact manifolds with corners. However, the submersion condition is not enough – for instance just take the identity map $[0, 1] \longrightarrow \mathbb{R}$ which has surjective differential but is not surjective. Insisting that a smooth map between manifolds with corners be surjective as well as have surjective differential at every point does lead to a fibration; it also ensures that the map be an interior b-map.

However, in the category of manifolds with corner there is a class of maps that is larger than this direct generalization of a fibration but which has enough of the properties to be very useful. It consists of the *b-fibrations*. To see where the defining conditions come from, recall that the differential f_* of a smooth map may be defined by duality from the pull-back. Namely the cotangent space of a manifold X at a point x is the quotient $T_x^*X = \mathcal{J}(x)/\mathcal{J}(x)^2$ of the ideal $\mathcal{J}(x) \subset \mathcal{C}^\infty(X)$ of smooth functions which vanish at x by the smaller ideal of functions which vanish to second order at x – which is spanned by the products of elements of $\mathcal{J}(x)$. Then $f^*\mathcal{J}(f(x)) \subset \mathcal{J}(x)$ and hence $f^* : T_{f(x)}^*X' \longrightarrow T_x^*X$ has dual which by definition is the differential $f_* : T_x X \longrightarrow T_{f(x)} X'$.

I have written out all this elementary stuff since on a manifold with corners there is a not-quite-obvious, but natural, generalization of it. First if ρ_1 and ρ_2 are defining functions for the same boundary hypersurface then $\rho_1 = a\rho_2$ where $0 < a \in \mathcal{C}^\infty(X)$. Thus $\log \rho_1 = \log \rho_2 + \log a$ where $\log a \in \mathcal{C}^\infty(X)$. It follows that the larger space of functions

$$(2.20) \quad \mathcal{C}_{\log}^\infty(X) = \{f : X \setminus \partial X \longrightarrow \mathbb{C}; f = \sum_j c_j \log \rho_j + f', f' \in \mathcal{C}^\infty(X), c_j \in \mathbb{C}\}$$

is independent of the boundary defining functions, ρ_j , used to define it and is therefore intrinsic. Moreover, interior b-maps define pull-back operations on it since under an such a map, see (2.5),

$$(2.21) \quad f^* \log(\rho'_i) = \sum_i \alpha_{ij} \log \rho_j + \log(a_i).$$

In local admissible coordinates $x_i = \rho_i$, the differentials of these functions are locally of the form

$$(2.22) \quad \sum_i (c_i + x_i u_i) \frac{dx_i}{x_i} + \sum_j v_j dy_j$$

for smooth functions u_i, v_j . Evaluating the coefficients at a point, i.e. taking the quotient, gives vector spaces ${}^b T_x^* X$ which are therefore naturally defined and combine to give a smooth vector bundle ${}^b T^* X$. The dual bundle, ${}^b T X$, is the one that the tangent vector fields, spanned locally by $x_i \partial_{x_i}$ and ∂_{y_j} form *all* the smooth sections of

$$(2.23) \quad \mathcal{V}_b(X) = \mathcal{C}^\infty(X; {}^b T X).$$

With this alternative tangent bundle in mind the b-differential is well-defined for any interior b-map by duality. What it does is tell us how the tangent vector fields behave under f ; at a boundary point it has a little more information in it than the usual differential. I will still denote it f_* since you can tell the difference since this $f_* : {}^bT_x X \rightarrow {}^bT_{f(x)} X'$.

Now, with this preamble it is not surprising that we define a *b-submersion* to be an interior b-map which has everywhere surjective b-differential. It is not quite clear that this condition is satisfied by fibrations in the category of manifolds with corners; it is but it is satisfied by other maps too. In particular

PROPOSITION 6. *The blow-down map $\beta : [M; F] \rightarrow M$ corresponding to blow up of any boundary face of a manifold with corners is a b-submersion.*

Blow maps for interior p-submanifolds, or any non-boundary face, are not b-submersions.

This is quite a useful concept but is not very close to that of a fibration. To get what we want, we need to impose another condition as well. This can be seen in various ways but the simplest is a global condition. Namely an interior b-map is said to be *b-normal* if no boundary hypersurface is mapped under it into a boundary face of codimension two (or higher of course). In terms of (2.5) this means that for each j there is at most one i such that $\alpha_{ij} \neq 0$. Indeed, $\{\rho_j = 0\}$ is mapped into $\{\rho'_i = 0\}$ under f if $\alpha_{ij} \neq 0$. Again a fibration is automatically b-normal, but a blow-down map (at least a non-trivial one, for a boundary face of codimension 2 or higher) is not b-normal.

DEFINITION 2. An interior b-map is a b-fibration if it is both a b-submersion and is b-normal.

It might be an interesting result if this condition implied that f was a fibration, but the truth is more interesting, namely it does not. To see a non-trivial example of a b-fibration consider the composite map of a blow-down and projection

$$(2.24) \quad f : [[0, 1]^2; \{0\}] \xrightarrow{\beta} [0, 1]^2 \xrightarrow{\pi_1} [0, 1].$$

Both maps are b-submersions, hence so is the composite which is clearly an interior b-map. Since the target manifold is a manifold with boundary, and hence has no boundary faces of codimension 2 or higher, the b-normality condition is automatically satisfied.

So, the claim I want to emphasize here is that b-fibrations are the replacements for fibrations in the category of manifolds with corners. I will try to justify this in various ways in the sequel. For the moment let me incorporate it into a definition. As I will explain below this definition needs to be expanded – here we are only considering the way smooth vector fields can degenerate at the boundary.

DEFINITION 3. Let $\mathcal{V} \subset \mathcal{V}_b(X)$ be a Lie algebra of smooth vector fields on a compact manifold with corners and suppose that \mathcal{V} contains all the smooth vector fields of compact support in the interior. A resolution of \mathcal{V} consists of a manifold with corners $X_{\mathcal{V}}^2$ which is obtained from X^2 by iterated blow up of p-submanifolds of the boundary (meaning at each stage – they do not have to be lifts of manifolds from X^2), so there is an overall blow-down map

$$(2.25) \quad \beta_{\mathcal{V}}^2 : X_{\mathcal{V}}^2 \rightarrow X^2$$

which is an identification of the interiors. It is further required that

- The factor exchange map lifts (extends by continuity from the interior) to be a smooth involution on $X_{\mathcal{V}}^2$.
- The diagonal lifts (to the closure of its intersection with the interior) to be a p-submanifold $\text{Diag}_{\mathcal{V}} \subset X_{\mathcal{V}}^2$.
- The elements of \mathcal{V} acting on the left factor lift (extend by continuity from the interior) to be smooth on $X_{\mathcal{V}}^2$ and to be collectively transversal to $\text{Diag}_{\mathcal{V}}$ at each point (i.e. they span the normal bundle to $\text{Diag}_{\mathcal{V}}$).
- The composite map $\pi_{L,\mathcal{V}} = \pi_L \circ \beta : X_{\mathcal{V}}^2 \rightarrow X$ is a b-fibration which is transversal to the lifted diagonal.

The first condition means that the last condition holds for the corresponding right stretched projection and similarly the third condition holds for the lift of the vector fields from the right factor.

These properties are enough to allow one to ‘microlocalize’ the Lie algebra to a ‘small’ space of pseudodifferential operators and to a ‘large’ space of pseudodifferential operators. Further properties (discussed below) ensure that the first is an algebra and in the second composition is possible under natural growth constraints. There are plenty of Lie algebras which cannot be resolved in this way (and also as we shall see there are more general notions of resolution if the conditions at the beginning that the vector fields be smooth and be arbitrary in the interior is dropped). Still there are lots of known examples.

PROBLEM 1 (Resolution problem). Is it possible to give a direct characterization of which Lie algebras are resolvable in this way?

PROBLEM 2 (Uniqueness problem). It is easy to see that $X_{\mathcal{V}}^2$ with the properties listed need not be unique. However, there should be some sort of uniqueness condition, meaning different resolutions should be closely related.

As noted above such a resolution is enough to define a space of operators. To prove composition results it is very convenient to go one step further and define a corresponding triple space.

DEFINITION 4. A *triple resolution* of X^3 associated to a resolution of a Lie algebra \mathcal{V} as in Definition 3 is a manifold with corners $X_{\mathcal{V}}^3$ obtained by iterated blow up of boundary p-submanifolds from X^3 , with overall blow-down map $\beta^3 : X_{\mathcal{V}}^3 \leftarrow X^3$, in such a way that

- The three factor exchange maps lift to diffeomorphisms
- The projection $\pi_F : X^3 \rightarrow X^2$ onto the right two factors lifts to a b-fibration $\pi_{F,\mathcal{V}} : X_{\mathcal{V}}^3 \rightarrow X_{\mathcal{V}}^2$ giving a commutative diagram

$$(2.26) \quad \begin{array}{ccc} X_{\mathcal{V}}^3 & \xrightarrow{\pi_{F,\mathcal{V}}} & X_{\mathcal{V}}^2 \\ \beta^3 \downarrow & & \downarrow \beta \\ X^3 & \xrightarrow{\pi_F} & X^2. \end{array}$$

Hence from the first condition the same is true of the other two projections π_S and π_C .

- The diagonal $\text{Diag}_{\mathcal{V}}$ in $X_{\mathcal{V}}^2$ lifts (to the closure of the inverse images of its interior) under each of the projections to three joint p-submanifolds which intersect precisely at the lift of the triple diagonal (which is therefore also a p-submanifold).

- The map $\pi_{F,\mathcal{V}}$ is transversal to the lifts of the diagonal under the other two projections.

The existence of such a triple resolution for \mathcal{V} guarantees the composition properties for operators mentioned above – these are made more precise later.

CONJECTURE 1. *There is always a triple resolution for any Lie algebra which has a resolution in the sense of Definition 3.*

In Definition 3 it was assumed that the initial object was a Lie algebra of smooth vector fields including all vector fields with compact support in the interior. This is rather an unreasonable restriction! I will include some examples below without giving a general *a priori* definition of resolution. The point is that both the single space, replacing X , and a replacement for the diagonal need to be chosen or constructed.

7. Examples of resolution of a vector fields

I did not cover these examples in the lectures at all, but I include here a substantial list (but by no means exhaustive) of Lie algebras which are known to have resolutions of this type introduced in Definition 3. Before doing so, let me give a result which reduces the workload a bit.

PROPOSITION 7. *If $\mathcal{V} \subset \mathcal{V}_b(X)$ is a Lie algebra of vector fields on a compact manifold with corners which has a resolution in the sense of Definition 3 then so does $\rho^\alpha \mathcal{V}$, with elements $\rho^\alpha V$, $V \in \mathcal{V}$, for any product of boundary defining functions ρ^α .*

Note that the Lie algebras obtained this way are by no means uninteresting and some are included in the list below.

CONJECTURE 2. *Let \mathcal{V} be a Lie algebra with a resolution as in Definition 3 and suppose $F \subset X$ is a boundary face of codimension 2 or greater. Then the Lie algebra $\mathcal{J}(F)\mathcal{V}$, formed by the span of the products of elements of \mathcal{V} and smooth function vanishing on F , has a resolution when lifted to $[X; F]$.*

(A)=b So, we start with a compact manifold with boundary X . The basic Lie algebra is $\mathcal{V}_b(X)$ itself. In boundary-adapted coordinates (which we always use) x and y_j it is spanned locally by

$$(2.27) \quad x\partial_x \text{ and } \partial_{y_j}, \quad j = 1, \dots, n-1, \quad n = \dim X.$$

It is resolved, as mentioned above, by blowing up the corner

$$(2.28) \quad X_b^2 = [X^2; (\partial X)^2] \text{ resolves } \mathcal{V}_b(X)$$

if ∂X is connected. If there are several components, $\partial X = \cup_j H_j$, of the boundary then there are different possible resolutions. The usual choice is just to take the products of the components of the boundary and consider

$$(2.29) \quad X_b^2 = [X^2; H_1 \times H_1; H_2 \times H_2; \dots; H_N \times H_N] \text{ resolves } \mathcal{V}_b(X).$$

One can also consider *all* the products between different boundary components. These products are disjoint in X^2 so the blow-up is independent of order

$$(2.30) \quad X_{ob}^2 = [X^2; \mathcal{M}_1(X) \times \mathcal{M}_1(X)] \text{ resolves } \mathcal{V}_b(X).$$

Here $\mathcal{M}_1(X)$ is the collection of boundary components; this is sometimes called the ‘overblown’ resolution.

The triple resolution associated to (2.29) is

$$(2.31) \quad X_b^3 = [X^3; (\partial X)^3; X \times (\partial X)^2; \partial X \times X \times \partial X; (\partial X)^2 \times X].$$

References:

- (B)=0 The next simplest case is the ‘zero Lie algebra’, $\mathcal{V}_0(X)$. (Other names have been used, especially in relation to conformal compactification because ‘zero’ seems to be interpreted as perjorative!) This consists of the smooth vector fields on X (in the usual sense) which vanish (hence the ‘zero’) at ∂X . It is spanned by

$$(2.32) \quad x\partial_x, x\partial_{y_j}.$$

Then

$$(2.33) \quad X_0^2 = [X^2; \partial \text{Diag}(X)] \text{ resolves } \mathcal{V}_0(X).$$

The associated triple resolution is analogous to (2.31)

$$(2.34) \quad X_0^3 = [X^3; \partial \text{Diag}_3; X \times \partial \text{Diag}; \dots]$$

where Diag_3 is the triple diagonal and the dots are the other boundaries of the other two partial diagonals – the images of the first one under the factor exchange maps.

References:

- (C)= ϕ -b More generally, and this is a construction we will apply several times below, we can consider a fibration of the boundary $\phi : \partial X \rightarrow B$. Then the fibred-boundary, also called ‘edge’ Lie algebra is

$$(2.35) \quad \mathcal{V}_{\phi\text{-b}}(X) = \{V \in \mathcal{V}_b(X); V \text{ is tangent to the fibres of } \phi\}.$$

We can now choose ‘boundary coordinates’ which are divided into two groups, z_l which are lifted from the base and y_j which induce coordinates on the fibres of ϕ . In terms of these the Lie algebra is spanned by

$$(2.36) \quad x\partial_x, x\partial_{y_k}, \partial_{z_l}$$

near the boundary. Within $\partial X \times \partial X$, the corner of X^2 , consider the fibre diagonal Diag_ϕ of ϕ . This is the set of pairs projecting to the same point in B , i.e. lying in the same fibre of ϕ . Then

$$(2.37) \quad X_\phi^2 = [X; \text{Diag}_\phi] \text{ resolves } \mathcal{V}_{\phi\text{-b}}(X).$$

In fact this includes the previous two cases as the special fibrations with one fibre (giving \mathcal{V}_b) and with point fibres (giving $\mathcal{V}_0(X)$). The triple resolution is given by the natural generalization

$$(2.38) \quad X_\phi^3 = [X^3; \text{Diag}_\phi^3; \pi_F^{-1}(\text{Diag}_\phi); \pi_S^{-1}(\text{Diag}_\phi); \pi_C^{-1}(\text{Diag}_\phi)].$$

References:-

- (D)=cu The next basic case is the cusp algebra $\mathcal{V}_{\text{cu}}(X)$. This is actually not well-defined but depends on the choice of some additional data. Namely one should fix a defining function for the boundary $x \in \mathcal{C}^\infty(X)$ up to a (positive) constant multiple and additional term $O(x^2)$. Geometrically this corresponds to an isomorphism of $N^*\partial X$ to $\partial X \times L$ for some real 1-dimensional vector space L . Different choices give different algebras but

they are identified by appropriate diffeomorphisms. Given the choice of defining function the cusp algebra

$$(2.39) \quad \mathcal{V}_{\text{cu}}(X) = \{V \in \mathcal{V}_{\text{b}}(X); Vx \in x^2\mathcal{C}^\infty(X)\}$$

is locally spanned by

$$(2.40) \quad x^2\partial_x, \partial_{y_j}.$$

Then

$$(2.41) \quad X_{\text{cu}}^2 = [X^2; \partial X \times \partial X; S] \text{ resolves } \mathcal{V}_{\text{cu}}(X)$$

where the first blow up gives X_{b}^2 and $S \subset \text{ff}(X_{\text{b}}^2)$ is a p-submanifold which can be defined as the flow-out of the lifted diagonal under the lift of elements of the cusp algebra. More usefully it can be written down as $s = 0$ where $s = (x - x')/(x + x')$ is a smooth function on X_{b}^2 obtained from the given defining function x on the left factor of X and x' on the right.

The triple resolution is now getting a little harder! We can start from X_{b}^3 in (2.31). Then we need to consider the three lifts of S from the three copies of X_{b}^2 and the corresponding triple submanifold T . The complexity comes from the fact that the inverse image of S under each of the stretched projections consists of *two* p-submanifolds, one in the ‘front face’ of X_{b}^3 (formed by the blow up of $(\partial X)^3$) and the other in the face coming from the corresponding corner of codimension two. Thus we have seven p-submanifolds to blow up T , three S_i^{ff} ’s and three S_i ’s and this is the order we need to use or we do not get a triple resolution

$$(2.42) \quad X_{\text{cu}}^3 = [X_{\text{b}}^3; T; S_i^{\text{ff}}, S_i].$$

Even a sketch of this is rather hard.

(E)= ϕ -cu The fibred-cusp algebras arising from a choice of cusp structure (actually less is needed, namely it is only needed ‘along the fibres’) and a fibration of the boundary ϕ as above:

$$(2.43) \quad \mathcal{V}_{\phi\text{-cu}}(X) = \{V \in \mathcal{V}_{\phi\text{-b}}(X); Vx \in x^2\mathcal{C}^\infty(X)\}.$$

It is locally spanned by

$$(2.44) \quad x^2\partial_x, x\partial_{z_i}, \partial_{y_j}$$

where the coordinates z are lifted from the base of the fibration on the boundary; it is resolved by a similar blow up to the cusp case:

$$(2.45) \quad X_{\phi\text{-cu}}^2 = [X_{\text{b}}^2; S_{\phi\text{-cu}}] \text{ resolves } \mathcal{V}_{\phi\text{-cu}}(X)$$

and a similar triple resolution, which I will not write down.

(F)=sc This is an extreme case of the previous example, where the fibration has points as fibres. It is spanned by

$$(2.46) \quad x^2\partial_x, x\partial_{y_j}.$$

I only mention it because it is important in applications. Note that it also follows from case (A) above and Proposition 7 that it has a resolution.

(G)=I- ϕ One can also iterate fibrations. That is, if one has a tower of fibrations

$$(2.47) \quad \partial X \xrightarrow{\phi_1} Y_1 \xrightarrow{\phi_2} Y_2 \xrightarrow{\phi_3} Y_3 \quad \cdots \xrightarrow{\phi_N} Y_N$$

then one can define a Lie algebra of vector fields with some higher jet information. Namly one can take a product decomposition of the manifold and extend the fibrations a little way out into the manifold so that the final base becomes $[0, \epsilon)_x \times Y_N$; denote these extended fibrations $\tilde{\phi}_j$. Then set

$$(2.48) \quad \mathcal{V}_{I-\phi} = \{V \in \mathcal{V}_b(X); V = V_1 + x^1 V_2 + x^2 V_3 + \cdots + x^{N-1} V_N + x^N V', \\ \text{where } V_j \text{ is tangent to the fibres of } \tilde{\phi}_j \text{ and } V' \in \mathcal{V}_b(X)\}.$$

Of course there are many ways to do the extension and the Lie algebra will depend on some of this information. There is a resolution using iterated blow ups and indeed a triple resolution.

(I)=b-H In fact it is not necessary to have a fibration of the boundary to produce an interesting Lie algebra. Suppose we simply have a subbundle $H \subset T\partial X$. Let $\alpha_i \in \mathcal{C}^\infty(X; \Lambda^1)$ be smooth 1-forms which define a lift of H from the boundary, in the sense that their joint null spaces at the boundary form a subbundle $\tilde{H} \subset T_{\partial X} X$ which is of rank one greater than H and for which $\tilde{H} \cap T\partial X = H$. Then we can set

$$(2.49) \quad \mathcal{V}_{b-H}(X) = \{V \in \mathcal{V}_0(X); \alpha_i(V) \in x^2 \mathcal{C}^\infty(X)\}.$$

This is a Lie algebra since

$$(2.50) \quad \alpha_i([V, W]) = V\alpha_i(W) - W\alpha_i(V) - d\alpha_i(V, W).$$

It is locally spanned by

$$(2.51) \quad x\partial_x, xV_l, x^2W_j$$

where the V_l restrict to the boundary to span H . Despite the notation the Lie algebra depends on more than H , rather on \tilde{H} . It has a resolution an a triple resolution.

(J)=ad The next example, the adiabatic algebra, is the first which does not satisfy the assumptions of Definition 3. It is fixed by a fibration, say of a compact manifold without boundary, $\phi: X \rightarrow Y$ with typical fibre Z . The vector fields we are interested in are on X but depend on a parameter, ϵ . For $\epsilon > 0$ they are just arbitrary vector fields depending smoothly on ϵ but at $\epsilon = 0$ we demand that they become tangent to the fibres of ϕ . Now, we can regard the parameter dependent vector fields as smooth vector fields on $\tilde{X} = X \times [0, 1]_\epsilon$ which satisfy

$$(2.52) \quad \mathcal{V}_{\text{ad}}(X) = \{V \in \mathcal{V}_b(\tilde{X}); V\epsilon = 0, V \text{ tangent to the fibres of } \phi \text{ at } \epsilon = 0\}.$$

In coordinates adapted to the fibration \mathcal{V}_{ad} is spanned by

$$(2.53) \quad \partial_{z_i}, \epsilon\partial_{y_j}.$$

Notice that the full space here is X_{ad} , not X , so $\mathcal{V}_{\text{ad}}(X_{\text{ad}})$ does not restrict to arbitrary vector fields in the interior – since there is no ϵ -derivative. Thus Definition 3 does not apply directly. Nevertheless there is a resolution in an essentially similar sense. The point however is that we do not need more than one ‘copy’ of the ϵ parameter, since it is a

parameter. The rôle of the diagonal is played by the fibre diagonal in ϵ . Thus the resolved space is

$$(2.54) \quad X_{\text{ad}}^2 = [X^2 \times [0, 1]; \text{Diag}(\phi) \times \{0\}]$$

where $\text{Diag}(\phi)$ is the fibre diagonal of ϕ . There are two maps back to the single space X_{ad} and all are b-fibrations with $\mathcal{V}_{\text{ad}}(X)$ lifting under each of them to be smooth and collectively transversal to the lifted fibre diagonal $\text{Diag}(X) \times [0, 1]$. The triple space follows the pattern that can be seen from the examples above.

References:-

(K)=b-cu Next consider an example of a ‘transition algebra.’ One such is the transition from $\mathcal{V}_{\text{b}}(X)$ to $\mathcal{V}_{\text{cu}}(X)$ for some compact manifold with boundary as a parameter ϵ approaches 0. Given the local bases (2.27) and (2.44) of these two algebras, the ‘obvious’ transition basis is

$$(2.55) \quad (x^2 + \epsilon^2)^{\frac{1}{2}} x \partial_x, \partial_{y_j}.$$

The first vector field is not smooth. It can be replaced by $(x + \epsilon)x\partial_x$ but this does not really mitigate the ‘lack of smoothness’. So the single space itself needs to be resolved

$$(2.56) \quad X_{\text{b-cu}} = [X \times [0, 1]; \partial \times \{0\}]$$

to which the vector fields in $\mathcal{V}_{\text{b-cu}}$ lift to be smooth. So in fact this is a setting rather similar to the preceding one and with a similar resolution:

$$(2.57) \quad X_{\text{b-cu}}^2 = [X^2 \times [0, 1]; (\partial X)^2 \times \{0\}; (\partial X)^2; S_{\text{b-cu}}; \partial X \times X \times \{0\}; X \times \partial X \times \{0\}].$$

Here $S_{\text{b-cu}}$ is a submanifold of the fact produced by the first blow up, of $(\partial X)^2 \times \{0\}$ which corresponds closely to S in the cusp case discussed above. There is a similar triple resolution.

References:-

(L)=cu- ϕ -cu Similar to the preceding case again but now a transition from cusp to fibred cusp, so the local spanning vector fields are

$$(2.58) \quad x^2 \partial_x, (x^2 + \epsilon^2)^{\frac{1}{2}} \partial_{y_j}, \partial_{z_i}$$

corresponding to a fibration of the boundary of a compact manifold with boundary as in case (E) above.

References:-

There are other such ‘transition algebras’.

(M)=b-f Now passing to rather more general cases, suppose X is a compact manifold with corners and $f : X \rightarrow X'$ is a b-fibration. Consider the space of smooth vector fields tangent to the fibres of f :

$$(2.59) \quad \mathcal{V}_{\text{b-f}}(X) = \{V \in \mathcal{V}_{\text{b}}(X); V f^* u = 0 \forall u \in \mathcal{C}^\infty(X')\}.$$

This has a resolution $X_{\text{b-f}}^2$ which is obtained from the fibre diagonal for f in X^2 by blow up in X^2 ; as usual there is a triple space.

Reference: None at the moment.

(N)=b-St The boundary stratification algebras. Consider an iterated conic space, a special type of stratified space. This is too hard to describe in a few sentences, but think of it as an iterated cone bundle. The space itself has a resolution to a compact manifold with corners X where the boundary hypersurfaces H_i are strictly order, corresponding to the ‘depth’ of the

stratum. Thus H_1 corresponds to the smallest singular stratum and H_N to the largest. Each of these H_j 's carries a fibration which 'remembers' the original stratum – thus its base is a resolution of the corresponding stratum. There are compatibility conditions for the strata at the corners – the leaves decrease as i increases. Here we are interested in the 'finite length' vector fields on the original manifold – which do not form a Lie algebra. To make them all smooth they are multiplied by a defining function for each boundary hypersurface of X . With arbitrary smooth coefficients the resulting vector fields on X form a Lie algebra which is an iterated version of the fibred boundary Lie algebra in (C). Near a point in the interior of H_i the Lie algebra reduces to the ϕ -b case. Near a corner, say of codimension 3, where the first three boundary hypersurfaces, H_1 , H_2 and H_3 meet, the algebra is spanned by

$$(2.60) \quad x_1 x_2 x_3 \partial_{x_1}, \quad x_2 x_3 \partial_{x_2}, \quad x_3 \partial_{x_3}, \quad x_1 x_2 x_3 \partial_{y'_i}, \quad x_2 x_3 \partial_{y''_i}, \quad x_3 \partial_{y'''_i}, \quad p a_{z_i}$$

where the tangential vector fields correspond to tangency to the different fibrations. Again this has a resolution, which I will not discuss.

(O)=m-sc There are various Lie algebras which correspond to the cases discussed above for a compact manifold with boundary but on a manifold with corners with no ordering of the boundary faces. For instance the basic Lie algebra $\mathcal{V}_b(X)$ on a manifold with corners, which is a special case of (M) where the b-fibration is the map to a point can be scaled as described in (7) by multiplying by a boundary defining function for each boundary hypersurface. This gives the multi-scattering algebra which is locally spanned in adapted coordinates by

$$(2.61) \quad \rho_{\text{tot}} x_1 \partial_{x_1}, \quad \rho_{\text{tot}} x_2 \partial_{x_2}, \quad \rho_{\text{tot}} \partial_{y_i} \rho_{\text{tot}} = x_1 \cdots x_n$$

near a corner of codimension two. The appropriate single space is X_{tot} obtained from X by blowing up all the boundary faces in order increasing with the dimension. There is a double and a triple resolution with which I will not bother you!

There are lots of other examples too. Some worked out in detail some not (yet).

8. Morse case again

9. b-calculus

10. Duality and distributions

11. Pull-back and push-forward

12. Smoothness under blow-up

13. Conormal distributions

14. Examples

15. More theorems!